



Productivity and Susceptibility Analysis for Selected California Fisheries

Report to

**California Ocean Science Trust
and
California Department of Fish and Wildlife**

December 2016

MRAG Americas, Inc.
www.mragamericas.com

Table of Contents

1	Introduction	1
1.1	Background	1
1.2	Productivity Susceptibility Analysis	2
2	Methodology	3
2.1	PSA Selection	3
2.2	Approach.....	7
2.3	Scoring	9
3	Results.....	12
4	Discussion	23
5	References	26
5.1	General References	26
5.2	Species Specific Data References	27

Project code:	US2212
Date of issue:	12/29/16
Prepared by:	Jill Swasey Dr. Erika Zollett Erin Wilson

1 Introduction

1.1 Background

In California, the Marine Life Management Act (MLMA) is the primary statute for management of marine fisheries. The Act, codified in 1999, provides guidelines for progressing toward ecosystem-based fisheries management which is comprehensive and proactive in order to achieve common objectives and meet identified standards. Since adoption of the MLMA, fisheries management has focused on targeted rulemakings and on the preparation of fishery management plans (FMPs) for select fisheries, sometimes in response to legislative action. FMPs can take a long time to prepare and can require extensive use of staff resources and funding; as a result, most of the state's fisheries have not yet benefited from this component of the MLMA but are nevertheless managed sustainably through the best efforts of the California Department of Fish and Wildlife (CDFW).

CDFW identified three needs to address the requirements of the MLMA for a large number of fisheries, given the resource limitations in terms of funds and staff:

1. A process for prioritizing future management actions both among and within fisheries;
2. A process for scaling those management actions to reflect the needs, risks, and values of each fishery together with the Department's capacity; and
3. A means of conveying up-to-date fisheries information in a way that's easy for stakeholders, researchers, and the public to navigate and digest.

To address the first need identified, the California Ocean Science Trust contracted MRAG Americas, Inc. to conduct a Productivity Susceptibility Analysis (PSA) on the state's most significant fisheries in terms of commercial value and recreational participation.¹ PSA is a method for assessing the vulnerability of a fishery species or stock, using a set of predetermined measureable attributes and scoring rankings. The approach assumes the level of vulnerability depends on two characteristics: the productivity of a species, which determines the rate at which it can sustain or recover from fishery-related impacts, and the susceptibility of the species or stock to fishing activities. As a result, fisheries management among various stocks can be prioritized in a transparent and consistent manner, based on their need for management action.

The PSA methodology is a powerful tool that allows stakeholders and regulators to gain perspective on the inherent risk of a fishery stock to fishing activities. It also allows scientists to clarify specifically where information is lacking and where to focus resources to collect more information, since attributes weigh differently on risk. This report focuses on the PSA analysis conducted by MRAG Americas, Inc. PSAs can be conducted alone or as part of a series of data analyses on vulnerability. PSAs do not consider stock status against approved biological reference points (although they do account for current biomass and fishing mortality levels), risk to the marine ecosystem, harvest guidelines for a stock, or effects of climate change. Using the information gained from a PSA, a potential secondary component

¹ The list fisheries for analysis was provided by CDFW and represent a diversity of stocks that span commercial and sport sectors, gear types, and coastal areas, and include finfish and invertebrates.

of the vulnerability analysis includes an Ecological Risk Assessment (ERA). An ERA assesses the risk a fishery poses to the ecosystem, and is a potential subsequent and separate phase of work.

The analysis provided here follows an established methodology in accordance with uses by NOAA Fisheries. PSAs provide a repeatable, expedient, and scientifically justifiable means to evaluate the relative vulnerability to fishery stocks for use in prioritizing fisheries for management action. They are primarily used for fishery stocks with moderate to strong long-term databases and knowledge of life history parameters, but this does not preclude stocks where less information is known from receiving management review. PSAs can provide managers with information needed to allocate resources appropriately in the short, medium, and long term. They are a first step in a tool kit of available approaches to assess and sustainably manage fisheries. They inform the users as to primary susceptibilities of the fishery stock and uncertainties in data gaps and quality of data used in the analysis. Built from expert opinion and best available scientific information, this type of risk-based approach allows managers the opportunity to decide the appropriate use of uncertainty and vulnerability in developing management strategies.

1.2 Productivity Susceptibility Analysis

Productivity Susceptibility Analysis (PSA) was originally developed to evaluate bycatch sustainability in the Australian prawn fishery by assessing productivity of bycatch stocks and their susceptibility to the fishery (Milton 2001, Stobutzki et al. 2001). In 2004, Australian Ecological Risk Assessment (AERA) team adapted the tool for a broader use in assessing vulnerability of an ecosystem (Hobday et al. 2004). Since then, it has been adapted for various assessments to evaluate vulnerability of ecosystems or stocks (Hobday et al. 2007; Rosenberg et al. 2007; Simpfendorfer et al. 2008; Patrick et al. 2009, 2010; Cope et al. 2011).

The outputs of a PSA vary, depending on the quality of data inputs and the methodologies applied. PSAs are useful for a baseline comparison among many species with varying levels of available information. For some stocks in California, stock assessments are available, some of which have been updated on a regular basis. For other stocks, little is known other than distribution or estimated life history characteristics, in addition to estimates of catch and fishing effort.

Using the PSA approach, productivity and susceptibility attributes of each stock (or assemblage) are examined and scored. The scores are then used in an equation to calculate overall vulnerability and are graphed to produce a PSA plot; the overall vulnerability scores and plot allow comparison of relative vulnerabilities with other units of analysis (target species, gear type, and fishing sector). Using this information, managers can prioritize stocks in high, medium, or low need of management attention and to identify areas where changes in management can most effectively reduce susceptibility. The PSA analysis also highlights gaps in understanding about a species' biology; improved information can allow for a better understanding of a stock's vulnerability.

2 Methodology

2.1 PSA Selection

Four versions of PSA methodologies were considered for use in this analysis; each was adapted from an approach developed by a joint Australian CSIRO/Australian Fisheries Management Authority project for Ecological Risk Assessment for the Effects of Fishing (ERAEF) (Hobday et al. 2007), and provide a good basis for a precautionary evaluation of vulnerability of fishery resources, including those modified by the National Oceanic and Atmospheric Administration (NOAA),² MRAG Americas,³ Marine Stewardship Council,⁴ and Monterey Bay Aquarium.⁵ Each method has slightly varying attributes and rankings.

In consultation with MRAG Americas and Ocean Science Trust, CDFW selected the NOAA's National Marine Fishery Service (NMFS) PSA approach for the analysis. The approach was developed as a means to evaluate the vulnerability of a stock in response to NMFS' revision of National Standard 1 (NS1) guidelines.

The NMFS approach was based on and developed from the attributes developed by Hobday et al. (2007). Several attributes that NMFS scientists perceived as redundant were removed. Retained attributes were those that were considered to be: (1) scientifically valid for calculating productivity or susceptibility of a stock, (2) useful at different scales (i.e., stocks of various sizes and spatial distributions), and (3) capable of being calculated for most fisheries (i.e., data availability). Attributes that were considered to have some but not all of these characteristics were retained, and four new attributes were added, including (1) recruitment pattern, (2) management strategy, (3) fishing rate relative to natural mortality, and (4) desirability/value of the fishery. The final NMFS PSA approach utilized 22 attributes (10 productivity, 12 susceptibility). Table 1 and Table 2 identify the final list of attributes, brief definitions and scoring criteria (the process of scoring is further detailed below). Additional details on each attribute are available in Patrick et al. (2009, 2010).

The NMFS PSA has been customized to specifically assess the vulnerability of U.S. fish stocks, based on definitions of becoming overfished or undergoing overfishing. An emphasis is placed on assessing data-poor stocks. The NMFS PSA was selected for this evaluation based on its inclusion of attributes that evaluate the management strategy and the value of a stock, and since it considers uncertainty in data sources. Missing data in the PSA are considered an endpoint in a continuum of data quality; therefore, it is suggested that managers account for data deficiencies and employ a precautionary approach when evaluating PSA results with limited or poor data.

² Patrick, W. S., P. Spencer, O. Ormseth, J. Cope, J. Field, D. Kobayashi, T. Gedamke, E. Cortés, K. Bigelow, W. Overholtz, J. Link, and P. Lawson. 2009. Use of productivity and susceptibility indices to determine stock vulnerability, with example applications to six U.S. fisheries. NOAA Tech. Memo. NMFSF/SPO-101. 90 pp.

http://www.nmfs.noaa.gov/sfa/laws_policies/national_standards/documents/patrick_2009_noaa_tech_memo_spo_101.pdf

³ MRAG Americas. 2009. Use of Productivity-Susceptibility Analysis (PSA) in Setting Annual Catch Limits for US Fisheries: An Overview. March 2009. http://www.mragamericas.com/wpcontent/uploads/2010/04/PSA_methodology.4.09.pdf

⁴ Marine Stewardship Council. 2014. MSC Fisheries Certification Requirements and Guidance V 2.0. <http://www.msc.org>

⁵ Monterey Bay Aquarium Seafood Watch Criteria for Fisheries (version March 31, 2014), PSA modified from MSC (2009).

Table 1. Productivity attributes and rankings used in this PSA analysis, derived from the NMFS PSA approach (adapted from Patrick et al. 2010).

Productivity Attribute	Definition	Ranking		
		High (3)	Moderate (2)	Low (1)
r	r is the intrinsic rate of population growth or maximum population growth that would occur in the absence of fishing at the lowest population size.	>0.5	0.5-0.16 (mid-point 0.10)	<0.16
Maximum Age	Maximum age is a direct indication of the natural mortality rate (M), where low levels of M are negatively correlated with high maximum ages.	< 10 years	10 - 30 years (mid-point 20)	> 30 years
Maximum Size	Maximum size is correlated with productivity, with large fish tending to have lower levels of productivity, although this relationship tends to degrade at higher taxonomic levels.	< 60 cm	60-150 cm (mid-point 105)	> 150 cm
von Bertalanffy Growth Coefficient (k)	The von Bertalanffy growth coefficient measures how rapidly a fish reaches its maximum size, where long-lived, low productivity stocks tend to have low values of k.	> 0.25	0.15-0.25 (mid-point 0.20)	< 0.15
Estimated Natural Mortality	Natural mortality rate directly reflects population productivity; stocks with high rates of natural mortality will require high levels of production in order to maintain population levels.	> 0.40	0.20-0.40 (mid-point 0.30)	< 0.20
Measured Fecundity	Fecundity (i.e., the number of eggs produced by a female for a given spawning event or period) is measured here at the age of first maturity.	> 10e4	10e2-10e3	< 10e2
Breeding Strategy	The breeding strategy of a stock provides an indication of the level of mortality that may be expected for the offspring in the first stages of life. Additional information in Winemiller 1989.	0	between 1 and 3	≥4
Recruitment Pattern	Stocks with sporadic and infrequent recruitment success often are long lived and thus may be expected to have lower levels of productivity.	highly frequent recruitment success (> 75% of year classes are successful)	moderately frequent recruitment success (between 10% and 75% of year classes are successful)	infrequent recruitment success (< 10% of year classes are successful)
Age at Maturity	Age at maturity tends to be positively related with maximum age (tmax); long-lived, lower productivity stocks will have higher ages at maturity than short-lived stocks.	< 2 years	2-4 years (mid-point 3.0)	> 4 years
Mean Trophic Level	The position of a stock within the larger fish community can be used to infer stock productivity; lower-trophic-level stocks generally are more productive than higher-trophic-level stocks.	<2.5	2.5-3.5 (mid-point 3)	>3.5

Table 2. Susceptibility attributes and rankings used in this PSA analysis, derived from the NMFS PSA approach (adapted from Patrick et al. 2010).

Susceptibility Attribute	Definition	Ranking		
		Low (1)	Moderate (2)	High (3)
Areal overlap	Areal overlap is the extent of geographic overlap between the known distribution of a stock and the distribution of the fishery.	< 25% of stock occurs in the area fished	Between 25% and 50% of the stock occurs in the area fished	> 50% of stock occurs in the area fished
Geographic concentration	Geographic concentration is the extent to which the stock is concentrated into small areas.	stock is distributed in > 50% of its total range	stock is distributed in 25% to 50% of its total range	stock is distributed in < 25% of its total range
Vertical overlap	Vertical overlap is the position of the stock within the water column (i.e., whether is demersal or pelagic) in relation to the fishing gear.	< 25% of stock occurs in the depths fished	Between 25% and 50% of the stock occurs in the depths fished	> 50% of stock occurs in the depths fished
Seasonal migrations	Seasonal migrations (i.e. spawning or feeding migrations) either to or from the fishery area could affect the overlap between the stock and the fishery.	Seasonal migrations decrease overlap with the fishery	Seasonal migrations do not substantially affect the overlap with the fishery	Seasonal migrations increase overlap with the fishery
Schooling, aggregation, and other behavioral responses	Behavioral responses of both individual fish and the stock respond to fishing activity.	Behavioral responses decrease the catchability of the gear	Behavioral responses do not substantially affect the catchability of the gear	Behavioral responses increase the catchability of the gear [i.e., hyperstability of CPUE with schooling behavior]
Morphological characteristics affecting capture	The ability of the fishing gear to capture fish varies based on their morphological characteristics (e.g., body shape, spiny versus soft rayed fins, etc.).	Species shows low selectivity to the fishing gear.	Species shows moderate selectivity to the fishing gear.	Species shows high selectivity to the fishing gear.
Desirability or value of the fishery	Highly valued fish stocks are assumed to be more susceptible to overfishing or to becoming overfished by recreational or commercial fishermen owing to increased effort.	Stock is not highly valued or desired by the fishery (<\$1/lb; <\$500K/yr landed; <33% retention).	Stock is moderately valued or desired by the fishery (\$1–\$2.25/lb; \$500K–\$10,000K/yr landed; 33–66% retention).	Stock is highly valued or desired by the fishery (>\$2.25/lb; >\$10,000K/yr landed; >66% retention).
Management strategy	The susceptibility of a stock to overfishing may largely depend on the effectiveness of fishery management procedures used to control catch.	Proactive management; sort requirements; individual specification; discard monitoring; biological data; representative fishery independent indices. Targeted stocks have catch limits and proactive accountability measures; Non-target stocks are closely monitored.	Reactive management; decent catch records; some assessment data; weak spatial knowledge; weakly informed indices. Targeted stocks have catch limits and reactive accountability measures	High catch uncertainty; low assessment data; no sorting; inadequate discard monitoring; low confidence in control rule. Targeted stocks do not have catch limits or accountability measures; Non-target stocks are not closely monitored.

Susceptibility Attribute	Definition	Ranking		
		Low (1)	Moderate (2)	High (3)
Fishing rate relative to M	As a conservative rule of thumb, it is recommended that M should be the upper limit of F so as to conserve the reproductive potential of a stock. (not avail without stock assessment)	<0.5	0.5 - 1.0	>1
Biomass of spawners (SSB) or other proxies	The extent to which fishing has depleted the biomass of a stock in relation to expected unfished levels offers information on realized susceptibility. This information is not available without a stock assessment.	B is > 40% of B0 (or maximum observed from time series of biomass estimates)	B is between 25% and 40% of B0 (or maximum observed from time series of biomass estimates)	B is < 25% of B0 (or maximum observed from time series of biomass estimates)
Survival after capture and release	Fish survival after capture and release varies by species, region, and gear type or even market conditions, and thus can affect the susceptibility of the stock.	Probability of survival > 67%	33% < probability of survival < 67%	Probability of survival < 33%
Impact of fisheries on EFH or habitat in general for non-targeted fish	A fishery may have an indirect effect on a species by adverse impacts on habitat.	Adverse effects absent, minimal, or temporary	Adverse effects more than minimal or temporary but are mitigated	Adverse effects more than minimal or temporary and are not mitigated

2.2 Approach

The PSA allows for the flexibility to define the unit of analysis. For this evaluation, the unit of analysis was defined as a combination of target species, gear, and fishing sector (commercial or sport). Some species were included in more than one unit of analysis since the gear type and/or sector differed enough to warrant additional analyses. CDFW provided the final list of species and sectors for evaluation, which included many of the state's most significant managed fisheries in terms of commercial value and recreational participation; the project scope required limiting to 45 units of analysis, which includes 21 finfish and 15 invertebrate species (Table 3).

PSA can be applied to single units of analysis or can consider cumulative impacts on a fishery, including the impacts of multiple gear types or sectors, bycatch, and takes throughout a species' range. This analysis evaluates only the susceptibility for fisheries in California and does not consider other fishing that may occur on a stock (e.g. for Survival After Capture and Release we assess the target species probability of survival once captured with the primary gear and not secondary sources of capture as bycatch in other fisheries). Users of this information should consider the possibility that fisheries in California may contribute a small amount of fishing pressure to a stock or species that is heavily fished in another state or country; this analysis would miss that cumulative pressure, thereby indicating a low risk for a high risk stock and potentially underestimating the overall vulnerability of the stock. Conversely, fishing activity in California may heavily exploit a small part of a non-overfished stock that has a wide range; the PSA would indicate a high risk for a low risk stock. Use of PSA results must consider the limitations of the selected approach for certain species.

Information and scores were initially generated by MRAG staff from readily available information. CDFW experts then reviewed available information and scoring to provide updated sources, results from Department research, expert opinion, and to either add scores not done by MRAG or change those that were based on preliminary or incomplete data by updating the attribute inputs and providing references. Experts involved in day-to-day management and data analysis for California fisheries possess knowledge that may not be contained in published literature. The information for the analyses and the corresponding scores were updated based on this expert evaluation. Once scores for productivity and susceptibility were determined, the overall vulnerability scores were calculated, using the following equation and only included attributes that could be scored: $v = \sqrt{(p - 3)^2 + (s - 1)^2}$.

Table 3. A unit of analysis is defined by the combination of target species, gear type, and fishing sector (commercial or sport). There are 45 units of analysis, which include 21 finfish and 15 invertebrate species. Some species are included in more than one unit of analysis. Hook and Line gear abbreviated as H&L.

Target Species	Gear Type	Fishery Sector	Target Species	Gear Type	Fishery Sector
Finfish			Invertebrates		
Barred sand bass	H&L	Sport	Bay shrimp	Beam trawl	Commercial
Barred surfperch	H&L	Sport	Brown rock crab	Trap	Commercial
Brown smoothhound shark	H&L	Sport	California spiny lobster	Hoop net	Sport
California barracuda	H&L	Sport	California spiny lobster	Trap	Commercial
California barracuda	H&L	Commercial	Dungeness crab	Trap	Sport
California corbina	H&L	Sport	Dungeness crab	Trap	Commercial
California halibut	H&L	Sport	Geoduck clam	Clam fork	Sport
California halibut	Trawl	Commercial	Giant red sea cucumber	Trawl	Commercial
California halibut	Gill net	Commercial	Kellett's whelk	Trap	Commercial
California halibut	H&L	Commercial	Market squid	Purse seine	Commercial
California sheephead	H&L	Sport	Ocean (pink) shrimp	Trawl	Commercial
California sheephead	Trap	Commercial	Pismo clam	Clam fork	Sport
Jacksmelt	H&L	Commercial	Red abalone	Abalone iron	Sport
Kelp bass	H&L	Sport	Red sea urchin	Hand rake (divers)	Commercial
Night smelt	A-frame	Commercial	Ridgeback prawn	Trawl	Commercial
Ocean whitefish	H&L	Sport	Spot prawn	Trap	Commercial
Pacific angel shark	Gill net	Commercial	Warty sea cucumber	Hand (divers)	Commercial
Pacific bonito	H&L	Sport			
Pacific bonito	H&L	Commercial			
Pacific hagfish	Trap	Commercial			
Pacific herring	Gill net	Commercial			
Redtail surfperch	H&L	Commercial			
Shiner seaperch	Trap	Commercial			
Spotted sand bass	H&L	Sport			
White croaker	H&L	Sport			
White seabass	H&L	Sport			
White seabass	Gill net	Commercial			
White sturgeon	H&L	Sport			

2.3 Scoring

Defining Scores

Productivity and susceptibility attributes are each scored based on predefined scoring bins, as provided in Patrick et al. (2009, 2010). Briefly, collected information and expert opinion provide the data that are compared with the scoring bins to identify a productivity or susceptibility score. All scores range from 1-3, but there is an inverse risk relationship between productivity and susceptibility (Figure 1). A fishery with low productivity and high susceptibility is more vulnerable to fishing activities than a fishery stock that is highly productive and/or has low susceptibility; a highly productive fishery may be able to recover from depletion or other impacts more quickly and is more likely to have a lower susceptibility.

Productivity is based on life history information. Susceptibility, however, evaluates the vulnerability of a stock to a given fishery; examining the susceptibility of a stock may present opportunity to reduce the risk. While scores range from 1-3, the NMFS PSA allows the flexibility to input intermediate scores (e.g. 1.5, 2.5) if deemed appropriate. The productivity (P) and susceptibility (S) attribute scores are averaged to yield separate, overall P and S scores.




Figure 1. Inverse relationship between productivity and susceptibility in measuring risk.

Treatment of Missing Data

Data were not available for every attribute for every unit of analysis. In some cases, expert opinion could be used to determine an appropriate score based on type of species or similar species where information was available, or based on the expert's best estimate. Where data were absent and expert opinion uncertain, select attributes not scored were left blank. As a precautionary measure, some ecological risk assessment approaches provide higher-level risk scores when data are missing in an attempt to avoid incorrectly identifying a high-risk stock as a low-risk. While precautionary, it may also confound the issue with data quality, where a data-poor stock could receive a high-risk result either from an abundance of missing data or from the analysis with available data. The approach utilized here isolates the treatment of uncertainty from the relative vulnerability and within the larger context of the quality of data used. In doing so, the approach chooses to decouple vulnerability and data quality by not

scoring attributes for which we had no information and report the overall quality of data separate from relative vulnerability.

Data Quality Index

Missing attribute scores were not factored into the overall scores but are reflected in the data quality score. Ranging from best data to no data, each attribute is assigned a data quality index to provide an estimate of information uncertainty. Scoring the quality of the data provides an additional lens through which the results should be considered. A data quality score of 1-5 was assigned to each attribute score, based on the reviewer's confidence in the data used where '1' reflects best available data and '5' indicates the absence of data and no attribute score (Table 4). Aggregate data quality scores for productivity and susceptibility data inputs are averaged from the individual attribute data quality scores; we can derive an overall data quality score from the average of the susceptibility and productivity data quality scores. The data quality score can be improved as more information becomes available on a fishery stock. The addition of information will reduce the uncertainty in the analysis but may not necessarily reduce the relative vulnerability. For invertebrates in particular, there were gaps in knowledge for certain attributes. However, certain attributes could be scored as high productivity, based on general species knowledge.

Incorporating data quality into the analysis allows poorly scored stocks to be flagged as either needing review of the scoring or indicating information is generally lacking for that stock. Therefore, the relative vulnerability scores that result from the analysis can be considered our best estimates based on best available scientific information, while the data quality index measures the information content in that best estimate (Patrick et al. 2009; Cope et al. 2011).

Table 4. The five tiers of data quality used when evaluating the productivity and susceptibility of an individual stock (adapted from Patrick et al. 2010).

Data Quality Tier	Description	Example
1	Best data. Information is based on collected data for the stock and area of interest and is established and substantial.	Data-rich stock assessment; published literature documenting methods used
2	Adequate data. Information is based on limited coverage and corroboration, or for some other reason is deemed not as reliable as tier-1 data.	Limited temporal or spatial data; relatively old information
3	Limited data. Estimates have high variation and limited confidence and may be based on studies of similar taxa or life history strategies.	Similar genus or family, etc.
4	Very limited data. Information is based on expert opinion or on general literature reviews from a wide range of species, or outside of region.	General data not referenced
5	No data. When there are no data on which to make even an expert opinion, the person using the PSA should give this attribute a “data quality” score of 5 and not provide a “productivity” or “susceptibility” score so as not to bias those index scores. When plotted, the susceptibility or productivity index score will be based on one less attribute, and will be highlighted as such by its related quality score.	

Information Review

For some species, model-generated data in Fishbase⁶ were the only information that was available for productivity attributes. In lieu of better data, these data were utilized if cases where other information for a particular species was also missing. In these cases, a low confidence data quality score was also given. The attributes with data gaps highlight uncertainty in scores to aid in interpreting overall vulnerability analyses, and where better data would provide a better evaluation of a stock’s vulnerability.

Nineteen different experts were consulted to review and aid in information review and scoring of the productivity and susceptibility attributes. Guidelines detailing the approach and explicit score rankings were provided to each expert for their review (see Table 1 and Table 2). Where scores based on the same or similar information varied considerably, a secondary review and decision was made prior to being finalized.

The NMFS PSA approach recognizes that not all of the productivity and susceptibility attributes will be equally useful for determining vulnerability. Previous versions of the PSA utilized an attribute weighting scheme in which higher weights were applied to the more important attributes as reflected through equations in vulnerability determinations (Stobutzki et al. 2001, Hobday et al. 2004, Rosenberg et al. 2007). The NMFS approach permits attribute weighting to customize the analysis. Attribute weighting is intended to reflect the relevance of the attributes in describing the productivity and susceptibility rather

⁶ An online global database of fish species (<http://www.fishbase.org/>).

than availability of data. No weights were applied to any attributes for any species analysis; given that our units of analysis include duplicative species from different fishery sectors we followed the NMFS recommendation by not assigning different weights among stocks within any given fishery.

3 Results

PSAs were conducted on the 45 units of analysis (Table 3), which included both commercial and sport sectors using a variety of gear types targeting finfish and invertebrate species. Where data or expert judgment provided, attributes were assigned a score; as described in Section 2.3, an accompanying data quality index was assigned to each score based on the information inputs, with the poorest data quality score assigned where an attribute score was missing. Overall data quality scores for productivity and susceptibility criteria (Figure 2) offer a snapshot of the general certainty in the data inputs. The majority of fisheries have good to moderate data quality; this should be considered in concert with the number of missing attribute scores. Further, certain attributes were scored more frequently than others. The frequency of productivity and susceptibility attributes scored and not scored is provided in Figure 3. A complete summary of the PSA results is provided in Table 5, this includes units of analysis, number of attributes scored, productivity (P), susceptibility (S), data quality (DQ) and vulnerability (V) scores. All data inputs and resources available were reviewed by CDFW staff to reduce uncertainty where expert opinion would apply and provide a transparent and consistent approach throughout.




Figure 2. Data quality plots for the productivity and susceptibility scores for all finfish and invertebrate fishery stocks analyzed demonstrates the distribution of data quality used across the analyses. Higher scores indicate poorer data quality (e.g., a score of 5 on either axis means the overall data quality is poorest and information is absent). Scores at the upper right corner therefore indicate the least-informed stocks. Scores closest to the origin are the most informed stocks. The vertical and horizontal lines provide a general guide to relative data quality, with values above 3 on either axis considered poorly informed scores.





Figure 3. The frequency of productivity attributes (top) and susceptibility attributes (bottom) scored for each of the forty-five units of analysis. Information derived from stock assessments were most frequently absent.

Table 5. Overall scores and results of the productivity–susceptibility analysis (PSA) for the 45 units of analysis. Abbreviations include: Commercial fishery (C), Sport fishery (S), Hook & Line gear (H&L), Productivity (P), Susceptibility (S), and Vulnerability (V). The number of attributes scored out of a possible 10 productivity and 12 susceptibility attributes. An * denotes indicator species that were selected by CDFW for multispecies fisheries. The scores are arranged first by increasing vulnerability and secondly grouped by species to allow sector comparisons. Degrees of vulnerability, as follows: lowest, $V < 1.8$; medium, $1.8 < V < 2.0$; high, $2.0 < V < 2.2$; and highest, $V > 2.2$.

Fishery	Species	Gear	No. of attributes scored		P Score	P Data Quality	S Score	S Data Quality	V
			P	S					
Jacksmelt (Silversides) (C)	<i>Atherinopsis californiensis</i>	H&L	7	9	2.43	2.60	1.67	2.67	0.88
Shiner seaperch (C) *	<i>Cymatogaster aggregata</i>	Trap	8	10	2.25	2.20	1.50	2.83	0.90
Dungeness Crab (S)	<i>Metacarcinus magister</i>	Trap	8	10	2.75	2.30	1.90	2.25	0.93
Dungeness Crab (C)	<i>Metacarcinus magister</i>	Trap	8	10	2.75	2.30	2.10	2.25	1.13
Night smelt (C) *	<i>Spirinchus starksi</i>	A-frame	8	10	2.56	2.50	1.85	2.33	0.96
Brown Rock Crab (C) *	<i>Cancer antennarius</i>	Trap	6	10	2.83	2.90	1.95	3.08	0.96
Pismo Clam (S)	<i>Tivela stultorum</i>	Clam Fork	7	10	2.43	2.60	1.80	2.50	0.98
White Croaker (S) *	<i>Genyonemus lineatus</i>	H&L	8	9	2.38	2.60	1.78	3.08	1.00
Bonito (C)	<i>Sarda chiliensis lineolata</i>	H&L	10	12	2.40	2.50	1.83	2.92	1.03
Bonito (S)	<i>Sarda chiliensis lineolata</i>	H&L	10	12	2.40	2.50	2.00	2.92	1.17
Pacific Hagfish (C)	<i>Eptatretus stoutii</i>	Trap	8	10	2.00	2.90	1.30	2.58	1.04
CA Corbina (S)	<i>Menticirrhus undulatus</i>	H&L	10	11	2.40	2.20	1.86	2.58	1.05
Ridgeback Prawn (C)	<i>Sicyonia ingentis</i>	Trawl	6	12	2.67	3.10	2.04	3.17	1.09
Redtail Surfperch (C) *	<i>Amphistichus rhodotus</i>	H&L	10	10	2.25	2.40	1.80	3.17	1.10
Kellett's Whelk (C)	<i>Kelletia kelletii</i>	Trap	6	8	2.33	3.60	1.94	2.75	1.15
Red Sea Urchin (C) *	<i>Strongylocentrotus franciscanus</i>	Hand rake	7	11	2.64	2.80	2.14	2.17	1.19
Spot Prawn (C)	<i>Pandalus platyceros</i>	Trap	6	10	2.50	3.20	2.10	2.58	1.21
Warty Sea Cucumber (C)	<i>Parastichopus parvimensis</i>	Diver	5	9	2.20	3.70	1.94	2.33	1.24
Barred surfperch (S) *	<i>Amphistichus argenteus</i>	H&L	10	11	2.00	2.40	1.77	2.67	1.26
White Seabass (S)	<i>Atractoscion nobilis</i>	H&L	9	11	2.11	1.90	1.91	1.83	1.27
CA Bay Shrimp (C)	<i>Crangon franciscorum</i>	Beam Trawl	6	9	2.33	3.30	2.11	3.08	1.30
Market Squid (C)	<i>Loligo (Doryteuthis) opalescens</i>	Purse Seine	9	11	2.50	2.50	2.23	3.25	1.33
CA Halibut (S)	<i>Paralichthys californicus</i>	H&L	10	12	1.90	2.00	1.75	3.00	1.33
CA Halibut (C)	<i>Paralichthys californicus</i>	H&L	10	12	1.90	2.00	1.75	3.00	1.33

Fishery	Species	Gear	No. of attributes scored		P Score	P Data Quality	S Score	S Data Quality	V
			P	S					
CA Halibut (C)	<i>Paralichthys californicus</i>	Trawl	10	12	1.90	2.00	2.08	3.00	1.54
CA Halibut (C)	<i>Paralichthys californicus</i>	Gillnet	10	12	1.90	2.00	2.13	3.00	1.57
Geoduck Clam (S)	<i>Panopea generosa</i>	Clam fork	10	11	1.90	2.10	1.77	2.67	1.34
CA Barracuda (C)	<i>Sphyraena argentea</i>	H&L	10	12	1.95	1.90	1.92	3.25	1.39
CA Barracuda (S)	<i>Sphyraena argentea</i>	H&L	10	12	1.95	1.90	2.00	3.25	1.45
Pink Shrimp (C)	<i>Pandalus jordani</i>	Trawl	7	9	2.57	2.80	2.33	2.75	1.40
Red Abalone (S)	<i>Haliotis rufescens</i>	Abalone Iron	9	12	1.89	1.60	1.88	2.00	1.41
White Seabass (C)	<i>Atractoscion nobilis</i>	Gillnet	9	11	2.11	1.90	2.14	1.83	1.44
Pacific Herring (C)	<i>Clupea pallasii</i>	Gillnet	10	11	2.35	2.10	2.32	2.50	1.47
Spotted Sand Bass (S)	<i>Paralabrax maculatofasciatus</i>	H&L	10	10	1.90	2.00	2.00	2.58	1.49
Barred Sand Bass (S)	<i>Paralabrax nebulifer</i>	H&L	10	11	2.05	1.60	2.18	1.92	1.52
CA Sheephead (S)	<i>Semicossyphus pulcher</i>	H&L	10	10	1.95	2.50	2.10	2.88	1.52
Kelp Bass (S)	<i>Paralabrax clathratus</i>	H&L	10	11	1.70	1.50	1.91	1.92	1.59
CA Sheephead (C)	<i>Semicossyphus pulcher</i>	Trap	10	10	1.95	2.50	2.30	2.79	1.67
CA Spiny Lobster (S)	<i>Panulirus interruptus</i>	Hoop Net	9	12	1.89	2.00	2.08	2.25	1.55
CA Spiny Lobster (C)	<i>Panulirus interruptus</i>	Trap	9	12	1.89	2.00	2.25	2.25	1.67
White Sturgeon (S)	<i>Acipenser transmontanus</i>	H&L	9	11	1.56	1.90	1.82	2.17	1.66
Giant Red Sea Cucumber (C)	<i>Parastichopus californicus</i>	Trawl	6	7	2.00	3.50	2.36	3.50	1.69
Ocean Whitefish (S)	<i>Caulolatilus princeps</i>	H&L	9	9	1.67	2.10	2.06	3.17	1.70
Brown Smoothhound Shark (S) *	<i>Mustelus henlei</i>	H&L	9	10	1.50	2.55	1.94	2.92	1.77
Pacific Angel Shark (C) *	<i>Squatina californica</i>	Gillnet	8	9	1.25	3.00	2.00	3.50	2.02

In compiling results across a variety of species and fisheries, it is meaningful to view the range of overall scores along with how scores relatively compare between species and fisheries. Figure 4 and Figure 5 depict productivity and susceptibility scores, respectively, for finfish and invertebrate species. It is important to note that PSA binning for scores is discrete, not continuous as the colors may reflect. Attribute cut-off scores are as follows: high productivity =1, moderate = 2, and low = 3; low susceptibility =1, moderate = 2, and high = 3. The average of all productivity or susceptibility attributes scored provides the final scores, respectively.

Of the 21 finfish species evaluated, productivities ranged from near high to near low, with the majority around moderate. Of the fifteen invertebrate species, seven were assessed to have high productivity and the remaining eight spanned the range of moderate productivity. Measures of susceptibility are specific to the fishing activity; therefore in the following visualizations we provide the fishery and sector (and for CA halibut we indicate gear type since there are three commercial gear types evaluated). The majority of invertebrate fisheries measured within a fairly narrow band of moderate susceptibility.




Figure 4. Summarized productivity scores across finfish (A) and invertebrate species (B). While productivity scores are discretely measured as high productivity =1, moderate = 2, and low = 3; we provide a comparative visualization to demonstrate the range of scores.








Figure 5. Summarized susceptibility scores across finfish (A) and invertebrate (B) fisheries. Abbreviations include California (CA), commercial (C), sport (S) and hook and line (H&L). Gears provided for CA halibut where three commercial sectors are analyzed. While susceptibility scores are discretely measured as low susceptibility =1, moderate = 2, and high = 3; we provide a comparative visualization to demonstrate the range of overall susceptibility scores.

The axes on PSA charts are oriented to align with how productivity and susceptibility are scored, with the lower left representing lowest vulnerability. Productivity measures across the x-axis, with the origin set at '3' (high productivity = low vulnerability); susceptibility measures along the y-axis with the origin set at '1' (low susceptibility = low vulnerability). Lowest vulnerabilities are those data points that are closest to the origin of the chart; the farther from the origin a data point sits, the higher the vulnerability. Data points that sit on the right side of the chart area measure low productivities; data points that sit towards the top of the chart area measure high susceptibilities. Relative vulnerabilities (V) and related data quality are provided for finfish fisheries in Figure 6 and for invertebrate fisheries in Figure 7. The contour lines divide regions of equal risk and group units of similar risk level. Overall data quality of the information used is reflected in the shading of symbols in the charts.








Figure 6. Relative vulnerabilities for finfish species: (A) CA halibut, (B) sharks, (C) white seabass and nearshore finfish (CA sheephead and ocean whitefish), (D) Pacific hagfish, Pacific herring and white sturgeon, (E) seabasses (barred sand bass, kelp bass and spotted sand bass) and nearshore bottom finfish (CA corbina and white croaker), (F) surf perch (shiner seaperch, barred surfperch, redtail surfperch), silversides (jacksmelt) and true smelt (night smelt), and (G) other nearshore pelagic finfish (bonito and CA barracuda). Symbol shading reflects overall data quality (poor >3.5 (solid black); moderate 2.0–3.5 (gray); and good <2.0 (white)). Lightly shaded background circles provide context for all finfish fisheries analyzed. Contours delineate areas of relative vulnerability (V, i.e., distance from the origin), with stocks of higher vulnerability above the solid line ($V = 2.0$), those of lower vulnerability below the dotted line ($V = 1.8$). Note that the productivity axis is in descending value in

order to make the top right quadrant of the plot the area of greatest vulnerability (i.e., the lowest productivity and highest susceptibility). P and S scores must be considered in concert with data quality for an appropriate assessment of risk from fishing activities and opportunities for increased information to alter the results. Abbreviations include California (CA), commercial (C) fishery, sport (S) fishery, and hook and line (H & L) gear. An * denotes indicator species selected by CDFW for multispecies fisheries.




Figure 7. Relative vulnerabilities for invertebrate species: (A) shrimp and prawn, (B) sea cucumber, market squid and sea urchin, (C) red abalone, clam and whelk, and (D) spiny lobster and crab. Symbol shading reflects overall data quality (poor >3.5 (solid black); moderate 2.0–3.5 (gray); and good <2.0 (white)). Lightly shaded background circles provide context for all invertebrate fisheries analyzed. Contours delineate areas of relative vulnerability (V , i.e., distance from the origin), with stocks of higher vulnerability above the solid line ($V = 2.0$), those of lower vulnerability below the dotted line ($V = 1.8$). Note that the productivity axis is in descending value in order to make the top right quadrant of the plot the area of greatest vulnerability (i.e., the lowest productivity and highest susceptibility). P and S scores

must be considered in concert with data quality for an appropriate assessment of risk from fishing activities and opportunities for increased information to alter the results. Abbreviations include California (CA), commercial (C) fishery, sport (S) fishery, and hook and line (H & L) gear. An * denotes indicator species selected by CDFW for multispecies fisheries.

4 Discussion

Productivity and Susceptibility Analysis (PSA) is a useful tool for assessing vulnerability of stocks to fishing activities and addressing both data poor and data rich species within the same analysis. The output is straightforward and allows for relatively quick, easy, and cost effective comparison among a large number of stocks, even when life history or fishery information is sparse. Considering limited funding for fishery management and environmental review, identifying the highest risk fisheries offers an opportunity to expend limited funds on the highest needs. Sorting fisheries by risk remains one of the most useful applications of PSA.

This analysis has already taken this preliminary step of grouping species into prioritization categories by identifying their relative vulnerabilities. Initial prioritization of fisheries was undertaken in designating the list of species and fishery sectors for assessments (Table 1); the results of this prioritization indicate:

- Species with low vulnerability which may therefore be a lower priority for further evaluation of management attention or stock status evaluations,
- Species for which management decisions can be deferred, or
- High risk species that likely need to move on to subsequent evaluations with either data rich or data poor methodologies.

PSA results can be used and interpreted in several additional ways as an evaluation planning process, taking into consideration the available resources for the CDFW; the approach can provide information on stock needs in terms of management attention and data improvement. For high risk species (such as those that are highly desirable, vulnerable to fishing activities, highly exploited, exhibit behavioral characteristics (e.g. spawning aggregations) that increase their susceptibility) the PSA highlights the susceptibility attributes which can potentially be altered through management to reduce a species' vulnerability to an identified fishery. Comparing relative vulnerabilities across species can provide information to understand where management might be focused to offer the greatest benefits for at-risk stocks.

Many management systems regulate multispecies fisheries though measures often aimed at indicator species or a small group of commonly co-occurring species, where information is presumed sufficient for management needs. However, PSA analyses and results provide information with which to consider whether managing to the data moderate and rich species is the most appropriate approach. In some circumstances, managing species with similar vulnerability profiles or re-evaluating the indicator species may be useful considerations:

- Fishery stock complexes are a common mechanism employed to manage multispecies fisheries, though these complexes may be based on similar biological characteristics (i.e. species guilds) or co-occurrence, rather than susceptibility to fishing activities. Stocks with similar overall vulnerabilities and characteristics can be grouped into stock complexes, and management considerations can be assessed for the consolidated group. Such an approach would reduce the resource needs for considering management on a fishery by fishery basis and manage to the identified vulnerabilities rather than developing measures that otherwise may not be appropriate for many species in the complex.
- Indicator species in a multispecies fishery are commonly identified as those with sufficient information on which to base management decisions. Given resource constraints of management, an alternate approach would be relying on the results of the PSA and delegating the most vulnerable stock as the indicator species, where possible, as the basis for management; however, analysis should be conducted on all species within a unit to identify the most vulnerable stock(s). This strategy would ensure that management measures would be precautionary enough to protect even the most vulnerable stocks (Patrick et al. 2010). The Department of Fisheries, Western Australia assesses the status of key retained species using PSA, and applies management intervention to the whole resource based on the most vulnerable key species (Fletcher et al., 2010, 2012; DoF 2011).

In terms of data improvement, data quality scores help identify where the greatest uncertainties in information exist. Together, the overall vulnerability and data quality scores suggest where data improvements are needed to understand the stock and fishery and to improve quality of the PSA. It is important for resource managers to consider trade-offs that might exist in efforts to fill data gaps. Directed research could be utilized to fill gaps in understanding a species' productivity but that may be unlikely to alter productivity or overall vulnerability scores. Increased confidence in the impacts from fishing activities are most likely to alter the vulnerabilities, though increased information could result in a higher vulnerability score where certain attributes scores were previously underestimated (such as the estimated areal overlap between the stock and fishing activities compared with actual overlap from improved spatial data). Improving data can be especially useful in data-limited situations and in prioritizing stock assessments (Cope et al. 2011).

Several methodological aspects should be considered when interpreting PSA results. Bias may exist with experts scoring fisheries they manage; an expert may unknowingly underestimate susceptibility scores (e.g. for management strategy), reflecting the expert's perception that the species is well managed. In this case, experts are most likely considering the susceptibility attributes (such as management strategy) of a fishery relative to other California fisheries with which they are familiar rather than to the full suite of management options that are utilized nationally or worldwide. For example, most sport fisheries in California are managed through bag limits, which are a form of catch limit, as well as other measures in many cases, and may be coupled with spatial and temporal closures and even size limits. However, monitoring and enforcement in these fisheries is often inadequate compared with commercial fisheries (e.g. Pacific herring or West Coast groundfish) that employ catch limits with monitoring and accountability measures. Care should be taken to minimize the opportunity for biases to result in

susceptibility scores that suggest a fishery is less susceptible than is actually the case. In this analysis, we ensured a consistent and transparent approach through the collaborative review of information inputs and documentation of references. In any approach that evaluates risk over a variety of species, maintaining consistency throughout the process is a critical component in the treatment of the information and utility of the results.

The NMFS PSA approach does not score an attribute with missing data; this is a considerable difference from some other PSA methodologies that default to a high risk score for attributes with missing data as discussed in Section 2.3. An approach that results in lower risk scores may provide less incentive to improve data quality or to reduce vulnerability by improving management. So while the NMFS PSA approach does not equate higher uncertainty with higher risk, which may result in lower vulnerability scores for data-poor species than other PSA approaches, the approach chooses to decouple the treatment of uncertainty from vulnerability allowing managers to decide where to add precaution. This analysis did not compare the results of the NMFS approach with other PSA approaches that default to a high risk score in absence of data; such an evaluation could produce interesting results to compare and consider. It is important to take these results in concert with the suite of attribute scores to isolate those attributes missing information. Additionally important, the PSA depends on how scoring bins are defined as low, moderate, and high for each attribute. These cut-off bins differ slightly among the PSA approaches, though it is unclear whether one approach is more or less conservative than other approaches.

Future steps towards understanding stock vulnerabilities in California fisheries could include expansions of the PSAs. Cumulative PSAs can be applied to look at certain impacts on a species such as various gear types or sectors, bycatch, and takes throughout a species' range. This approach would consider the full spatial range of all fishing activities and related impacts on the stock, and would be the most comprehensive assessment of a stock's vulnerability; however, collectively accounting for all of the variables requires additional resources and there are likely to be larger information gaps across the range of impacts on a stock. The approach employed here is consistent with the NMFS methodology and allows comparison between individual units of assessment and consideration of differences between sectors or gears. There is the risk of underestimating vulnerability by not accounting for cumulative impacts such as full range and overlap of multiple fishing pressures. PSA results can also be applied towards subsequent phases of work including an Ecological Risk Assessment (ERA), which assesses the risk a fishery poses to the ecosystem. The most appropriate next step will depend on needs identified by the CDFW.

5 References

5.1 General References

- Cope, J.M., J. DeVore, E. J. Dick, K. Ames, J. Budrick, D.L. Erickson, J. Grebel, G. Hanshew, R. Jones, L. Mattes, C. Niles, and S. Williams. 2011. An Approach to Defining Stock Complexes for U.S. West Coast Groundfishes Using Vulnerabilities and Ecological Distributions. *North American Journal of Fisheries Management* 31 (4): 589-604.
- Department of Fisheries (DoF). 2011. Resource Assessment Framework for Finfish Resources in Western Australia. Fisheries Occasional Publication No. 85. Department of Fisheries, Western Australia. 24 pp. http://www.fish.wa.gov.au/Documents/occasional_publications/fop085.pdf
- Fletcher, W.J., J. Shaw, S.J. Metcalf, and D.J. Gaughan. 2010. An ecosystem based fisheries management framework: the efficient, regional level planning tool for management agencies. *Mar. Policy* 34: 1226–1238.
- Fletcher, W.J., D.J. Gaughan, S.J. Metcalfe, and J. Shaw. 2012. Using a regional level, risk based framework to cost effectively implement Ecosystem based Fisheries Management (EBFM). In Kruse, G.H., H.I. Browman, K.I. Cochrane, D. Evans, G.S. Jamieson, and P.A. Livingstone (Eds.), *Global Progress on Ecosystem Based Fisheries Management*, Alaska Sea Grant College Program. pp. 129–146
10.4027/gpebfm.2012.07
- Hobday, A. J., A. Smith, and I. Stobutzki. 2004. Ecological risk assessment for Australian Commonwealth fisheries, final report. Report R01/0934 to the Australian Fisheries Management Authority, Canberra, Australia. 72 pp. http://www.afma.gov.au/research/reports/2004/r01_0934.pdf
- Hobday, A. J., A. Smith, H. Webb, R. Daley, S. Wayte, C. Bulman, J. Dowdney, A. Williams, M. Sporicic, J. Dambacher, M. Fuller, and T. Walker. 2007. Ecological Risk Assessment for the Effects of Fishing: Methodology. Report R04/1072 for the Australian Fisheries Management Authority, Canberra.
- Milton, D. A. 2001. Assessing the susceptibility to fishing of populations of rare trawl bycatch: sea snakes caught by Australia's Northern Prawn Fishery. *Biol. Conserv.* 101: 281–290.
- Patrick, W. S., P. Spencer, O. Ormseth, J. Cope, J. Field, D. Kobayashi, T. Gedamke, E. Cortés, K. Bigelow, W. Overholtz, J. Link, and P. Lawson. 2009. Use of productivity and susceptibility indices to determine stock vulnerability, with example applications to six U.S. fisheries. NOAA Tech. Memo. NMFSF/SPO-101. 90 pp. http://www.nmfs.noaa.gov/sfa/laws_policies/national_standards/documents/patrick_2009_noaa_tech_memo_spo_101.pdf
- Patrick, W. S., P. Spencer, J. Link, J. Cope, J. Field, D. Kobayashi, P. Lawson, T. Gedamke, E. Cortes, O. Ormseth, K. Bigelow, and W. Overholtz. 2010. Using productivity and susceptibility indices to assess the vulnerability of United States fish stocks to overfishing. *U.S. National Marine Fisheries Service Fishery Bulletin* 108: 305–322.

http://www.nmfs.noaa.gov/sfa/laws_policies/national_standards/documents/patrick_et_al_2010_psa.pdf

Rosenberg, A, D. Agnew, E. Babcock, A. Cooper, C. Mogensen, R. O'Boyle, J. Powers, G. Stefansson, and J. Swasey. 2007. Setting annual catch limits for U.S. fisheries: An expert working group report. MRAG Americas, Washington, D.C. 36 pp. https://www.mragamericas.com/wp-content/uploads/2010/04/PSA_Workshop-Report_May-09_MRAG-FINAL.pdf

Simpfendorfer, C., E. Cortés, M. Heupel, E. Brooks, E. Babcock, J. K. Baum, R. McAuley, S. F. J. Dudley, J. D. Stevens, S. Fordham, and A. Soldo. 2008. An integrated approach to determining the risk of over-exploitation for data-poor pelagic Atlantic sharks. An Expert Working Group Report, Lenfest Ocean Program, Washington, D.C. 22 pp.

Stobutzki, I., M. Miller, and D. Brewer. 2001. Sustainability of fishery bycatch: a process for assessing highly diverse and numerous bycatch. *Environmental Conservation* 28: 167-181.

Winemiller, K. O. 1989. Patterns of variation in life history among South American fishes in seasonal environments. *Oecologia* 81: 225-241.

5.2 Species Specific Data References

5.2.1 Barracuda

Bottinelli, D.J. and L.G. Allen. 2007. A Re-evaluation of age, growth, and batch fecundity in the California Barracuda, *Sphyræna argentea*, from southern California based on specimens taken from 2000 to 2002. *California Fish and Game* 93(4): 167-199.

California Department of Fish and Game (CDFG). 2001. California's Living Marine Resources: A Status Report. The Resources Agency, The California Department of Fish and Game. December 2001. 594 pp. <https://www.wildlife.ca.gov/Conservation/Marine/Status/2001>

California Department of Fish and Wildlife (CDFW). 2013. California Marine Sportfish Identification: Other Fishes. California Department of Fish and Wildlife Marine Region. <https://www.wildlife.ca.gov/Fishing/Ocean/Fish-ID/Sportfish/Other-Fishes#barracuda>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife. <https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

Eschmeyer, W.N., E.S. Herald, and H. Hamman, 1983. A field guide to Pacific coast fishes of North America. Houghton Mifflin Company, Boston, U.S.A. 336 pp.

Fishbase. *Sphyræna argentea*, Pacific barracuda. <http://www.fishbase.org/summary/3678>

IGFA. 2001. Database of IGFA angling records until 2001. IGFA, Fort Lauderdale, USA.

MRAG Americas, Inc. 2016. Assessment of CDFW Data Collections, Final Report of Phase 1 & 2. A Report for California Department of Fish and Wildlife and Resources Legacy Fund. MRAG Americas, Essex, MA.

Pinkas, L. 1966. A management study of the California barracuda, *Sphyræna argentea* Girard. California Department of Fish and Game, Fish Bulletin 134, 58 pp.

Radovich, J. 1961. Relationships of some marine organisms of the northeast Pacific to water temperatures particularly during 1957 through 1959. Calif. Dep. Fish Game Fish Bull. 112: 62 pp.

Shanks, A.L. and G.L. Eckert. 2005. Population persistence of California Current fishes and benthic crustaceans: a marine drift paradox. Ecol. Monogr. 75: 505-524.

5.2.2 Barred Sand Bass

CDFG. 2004. Annual Status of the Fisheries Report Through 2003. Report to the Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Game Marine Region, December 2004.

<https://www.wildlife.ca.gov/Conservation/Marine/Status#28027680-status-of-the-fisheries-report-through-2003>

CDFW. 2013. California Marine Sportfish Identification: Sea Bass. California Department of Fish and Wildlife Marine Region. <https://www.wildlife.ca.gov/Fishing/Ocean/Fish-ID/Sportfish/Sea-Bass#kelp>

CDFW. 2016. Saltwater Basses Fishery Information. California Department of Fish and Wildlife Marine Region. <https://www.wildlife.ca.gov/Conservation/Marine/SCFRMP/Saltwater-Bass#newregs>

CDFW. Unpublished data/ongoing study on growth. California Department of Fish and Wildlife Marine Region.

CDFW. Unpublished data/ongoing study on “hot spot” analyses. California Department of Fish and Wildlife Marine Region.

Fishbase. *Paralabrax nebulifer*, Barred sand bass.

<http://www.fishbase.org/Summary/speciesSummary.php?ID=3337&AT=barred+sand+bass>

IGFA. 2001. Database of IGFA angling records until 2001. IGFA, Fort Lauderdale, USA.

Jarvis, E.T., C. Linardich, and C.F.Valle. 2010. Spawning-related movements of barred sand bass, *Paralabrax nebulifer*, in southern California: Interpretations from two decades of historical tag and recapture data. Southern California Academy of Sciences Bulletin, 109: 123-143.

<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=48389&inline>

Jarvis, E.T., H.L. Gliniak, and C.F.Valle. 2014. Effects of fishing and the environment on the long-term sustainability of the recreational saltwater bass fishery in southern California.

<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=90973&inline>

Love, M. S., A. Brooks, D. Busatto, J. Stephens, and P. A. Gregory. 1996. Aspects of the life histories of the kelp bass, *Paralabrax clathratus*, and barred sand bass, *P. nebulifer*, from the Southern California Bight. Fishery Bulletin 94: 472-481.

Shanks, A.L. and G.L. Eckert, 2005. Population persistence of California Current fishes and benthic crustaceans: a marine drift paradox. Ecol. Monogr. 75: 505-524.

Then, A.Y., J.M. Hoenig, N.G. Hall, and D.A. Hewitt. 2015. Evaluating the predictive performance of empirical estimators of natural mortality rate using information on over 200 fish species. ICES Journal of Marine Science 72(1): 82–92.

5.2.3 Barred Surfperch

Breder, C.M. and D.E. Rosen. 1966. Modes of reproduction in fishes. T.F.H. Publications, Neptune City, New Jersey. 941 pp.

Carlisle, J., J. Schott, and N. Abramson. 1960. The barred surfperch (*Amphistichus argenteus* Agassiz) in Southern California. Calif. Fish Game Fish. Bull. (109). 79 pp.

CDFW. 2013. Status of the Fisheries Report, an Update through 2011. Report to the California Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Wildlife Marine Region. January 2013.

<https://www.wildlife.ca.gov/Conservation/Marine/Status#28027677-status-of-the-fisheries-report-through-2011>

CDFW. Unpublished data on maximum size and growth. California Department of Fish and Wildlife Marine Region.

Eschmeyer, W.N., E.S. Herald, and H. Hammann. 1983. A field guide to Pacific coast fishes of North America. Houghton Mifflin Company, Boston, U.S.A. 336 pp.

Fishbase. *Amphistichus argenteus*, Barred surfperch.

<http://www.fishbase.org/Summary/speciesSummary.php?ID=3622&AT=barred+surfperch>

Pauly, D. 1978. A preliminary compilation of fish length growth parameters. Ber. Inst. Meereskd. Christian-Albrechts-Univ. Kiel (55): 1-200.

Shanks, A.L. and G.L. Eckert. 2005. Population persistence of California Current fishes and benthic crustaceans: a marine drift paradox. Ecol. Monogr. 75: 505-524.

5.2.4 Bay Shrimp

California Department of Fish and Game (CDFG). 2001. California's Living Marine Resources: A Status Report. The Resources Agency, The California Department of Fish and Game. December 2001.

<https://www.wildlife.ca.gov/Conservation/Marine/Status/2001>

CDFW. 2015. Commercial Digest California Fishing Regulations. 2015-2016. California Department of Fish and Wildlife Marine Region. <https://www.wildlife.ca.gov/Conservation/Marine/Groundfish/Commercial-Regulations>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife.
<https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

CDFW. 2016. Commercial landing receipts, Commercial Fishery Information System (CFIS) (Extracted and analyzed 7/2016).

Jennings, S., M. Kaiser, and J. Reynolds. 2001. Marine fisheries ecology. Blackwell Science Ltd., Malden, MA.

NMFS. 2005. Essential Fish Habitat Designation and Minimization of Adverse Impacts, Pacific Coast Groundfish Fishery Management Plan: Environmental Impact Statement.
http://www.westcoast.fisheries.noaa.gov/publications/nepa/groundfish/final_groundfish_efh_eis.html

Siegfried, C.A. 1989. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (Pacific Southwest) --crangonid shrimp. U.S. Fish Wildl. Serv. Biol. Rep. 82 (11.125). U.S. Army Corps of Engineers, TR EL-82-4. 18pp.

5.2.5 Bonito

Anonymous, 1994. U. S. GLOBEC - Global ocean ecosystems dynamics, a component of the U. S. Global Change Research Program. Eastern Boundary Current Program-A Science Plan for the California Current, Rep. No. 11, August.

Black, G. 1979. Maturity and spawning of the Pacific bonito, *Sarda chiliensis lineolata*, in the eastern north Pacific. Calif. Dept. of Fish and Game Marine Resources Technical Report No. 41. 61 p.

Campbell, G. and R.A. Collins. 1975. The age and growth of the Pacific bonito, *Sarda chiliensis*, in the eastern north Pacific. Calif. Fish and Game 61(4): 181-200

CDFG. 2010. California's Living Marine Resources: Status of the Fisheries Report through 2008. Report to the California Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Game Marine Region, August 2010.
<https://www.wildlife.ca.gov/Conservation/Marine/Status#28027678-status-of-the-fisheries-report-through-2008>

CDFW. 2009. California Marine Life Protection Act Initiative Methods Used to Evaluate Draft MPA Proposals in the MLPA South Coast Study Region (DRAFT) ADDITIONS TO Chapter 3.0 – Protection Levels, Revised March 23, 2009. California Department of Fish and Wildlife Marine Region.
https://www.dfg.ca.gov/marine/pdfs/agenda_040109d1.pdf

CDFW. 2015. California Marine Sportfish Identification: Tuna & Mackerels. California Department of Fish and Wildlife Marine Region. <https://www.wildlife.ca.gov/Fishing/Ocean/Fish-ID/Sportfish/Tuna-And-Mackerels#bonito>

Collette, B.B. and C.E. Nauen, 1983. FAO Species Catalogue. Vol. 2. Scombrids of the world. An annotated and illustrated catalogue of tunas, mackerels, bonitos and related species known to date. Rome: FAO. FAO Fish. Synop. 125(2). 137 pp.

Collins, R.A., D. Huppert, A. MacCall, J. Radovich, and G. Stauffer. 1980. Pacific Bonito Management Information Document. Marine Resources Technical Report No. 44, 94 p. California Dept. of Fish and Game.

FishBase. *Sarda chiliensis*, Eastern Pacific bonito.

<http://www.fishbase.org/Summary/SpeciesSummary.php?ID=113&AT=bonito>

MRAG Americas, Inc. 2016. Assessment of CDFW Data Collections, Final Report of Phase 1 & 2. A Report for California Department of Fish and Wildlife and Resources Legacy Fund. MRAG Americas, Essex, MA.

Snow, J. 2016. Mexico Fish, Flora, & Fauna. Pacific Bonito, *Sarda chiliensis*. <http://www.mexican-fish.com/pacific-bonito/>

5.2.6 Brown Rock Crab

Carroll, J.C., and R.N. Winn. 1989. Species Profiles: life histories and environmental requirements of coastal fishes and invertebrates (Pacific Southeast)--brown rock crab, red rock, and yellow crab. US Fish and Wildlife Service Biological Report. 82 (11.117). US Army Corps of Engineers, TR EL-82-4. 16 pp.

CDFG. 2004. Annual Status of the Fisheries Report Through 2003. Report to the Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Game Marine Region, December 2004.

<https://www.wildlife.ca.gov/Conservation/Marine/Status#28027680-status-of-the-fisheries-report-through-2003>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife.

<https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

MRAG Americas, Inc. 2016. Assessment of CDFW Data Collections - Final Report of Phase 1 & 2. A Report for California Department of Fish and Wildlife and Resources Legacy Fund May 3, 2016.

Safina Center for Seafood Solutions. 2014. Seafood Watch Report Pacific Rock Crabs: Brown, Red and Yellow (*Cancer antennarius*, *Cancer productus*, and *Cancer anthonyi*). May 12, 2014.

http://www.seafoodwatch.org/-/m/sfw/pdf/reports/c/mba_tsc_pacificrockcrabreport.pdf

5.2.7 Brown Smoothhound Shark

Breder, C.M. and D.E. Rosen. 1966. Modes of reproduction in fishes. T.F.H. Publications, Neptune City, New Jersey. 941 pp.

Cailliet, G.M., K.G. Yudin, S. Tanaka, and T. Taniuchi. 1990. Growth characteristics of two populations of *Mustelus manazo* from Japan based upon cross-readings of vertebral bands. NOAA Tech. Rep. NMFS 90: 167-176.

CDFW. 2013. California Marine Sportfish Identification, Sharks. California Department of Fish and Wildlife Marine Region. <https://www.wildlife.ca.gov/Fishing/Ocean/Fish-ID/Sportfish/Sharks>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife. <https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

CDFW. 2016. California Ocean Sport Fishing Regulations, 2016-2017. California Department of Fish and Wildlife Marine Region. <https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=116558>

Chabot, C.L. and B.M. Haggin. 2014. Frequency of multiple paternity varies between two populations of brown smoothhound shark, *Mustelus henlei*. Mar. Biol. 161: 797-804.

Compagno, L.J.V. 1984. Sharks of the World: an annotated and illustrated catalogue of the shark species known to date. Part 2. Carcharhiniformes. FAO, Rome.

Ebert, D.A. 2003. Sharks, Rays and Chimaeras of California. University of California Press, Berkley.

Eschmeyer, W.N., E.S. Herald, and H. Hammann. 1983. A field guide to Pacific coast fishes of North America. Houghton Mifflin Company, Boston, USA. 336 pp.

Fleming, K. 1999. Elasmobranchs. In: J. Orsi (ed.), Report on the 1980-1995 fish, shrimp, and crab sampling in the San Francisco Estuary, California. California Fish and Game.

Fishbase. *Mustelus henlei*, Brown smooth-hound.

<http://www.fishbase.org/Summary/speciesSummary.php?ID=2540&AT=brown+smoothhound>

Love, M.S., C.W. Mecklenburg, T.A. Mecklenburg, and L.K. Thorsteinson. 2005. Resource Inventory of Marine and Estuarine Fishes of the West Coast and Alaska: A Checklist of North Pacific and Arctic Ocean Species from Baja California to the Alaska-Yukon Border. U.S. Department of the Interior, U.S. Geological Survey, Biological Resources Division, Seattle, Washington, 98104.

MRAG Americas, Inc. 2016. Assessment of CDFW Data Collections - Final Report of Phase 1 & 2. A Report for California Department of Fish and Wildlife and Resources Legacy Fund. May 3, 2016.

Pérez-Jiménez, J.C. 2006. Biología y taxonomía de los tiburones del género *Mustelus* (Elasmobranchii) de la región norte del Golfo de California. PhD Thesis, CICESE.

Pérez-Jiménez, J.C. and A.B. Carlisle. 2009. *Mustelus henlei*. The IUCN Red List of Threatened Species 2009: e.T161648A5472414. <http://dx.doi.org/10.2305/IUCN.UK.2009-2.RLTS.T161648A5472414.en>.

Smith, S.W., D.W. Au, and C. Show. 1998. Intrinsic rebound potential of 26 species of Pacific sharks. Mar. Freshwat. Res. 49: 663-678.

Yudin, K.G. 1987. Age, Growth and Aspects of the Reproductive Biology of Two Sharks, The Gray Smoothhound (*Mustelus californicus*) and the Brown Smoothhound *M. Henlei* from Central California. M.A. Thesis, San Francisco State University.

Yudin, K.G. and G.M. Cailliet. 1990. Age and growth of the gray smoothhound, *Mustelus californicus*, and the brown smoothhound, *M. henlei*, from central California. *Copeia* 1990(1): 191-204.

5.2.8 California Corbina

CDFG. 2001. California's Living Marine Resources: A Status Report. The Resources Agency, The California Department of Fish and Game. December 2001. 594 pp.

<https://www.wildlife.ca.gov/Conservation/Marine/Status/2001>

CDFW. 2013. California Marine Sportfish Identification: Croakers. California Department of Fish and Wildlife Marine Region. <https://www.wildlife.ca.gov/Fishing/Ocean/Fish-ID/Sportfish/Croakers#corbina>

Chao, L.N. 1995. Sciaenidae. Corvinas, barbiches, bombaches, corvinatas, corvinetas, corvinillas, lambes, pescadillas, roncachos, verrugatos. p. 1427-1518. In W. Fischer, F. Krupp, W. Schneider, C. Sommer, K.E. Carpenter and V. Niem (eds.) *Guia FAO para identificacion de especies para los fines de la pesca. Pacifico Centro-oriental*. 3 volumes. 1813 pp.

Eschmeyer, W.N., E.S. Herald, and H. Hammann. 1983. *A field guide to Pacific coast fishes of North America*. Houghton Mifflin Company, Boston, U.S.A. 336 pp.

FishBase. *Menticirrhus undulatus*, California kingcroaker.

<http://www.fishbase.org/Summary/speciesSummary.php?ID=3587&AT=California+corbina>

Gliniak, H.G. 2013. California corbina, *Menticirrhus undulatus*. In: Larinto, T. (ed.). *Status of the Fisheries Report – An Update Through 2011*. CA Dept. of Fish and Wildlife.

<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=65551&inline>

Joseph, D. 1962. Growth characteristics of two southern California surf fishes, the California corbina and spoonfin croaker, family Sciaenidae. *Calif. Fish. Game, Fish. Bull.* (119): 1-54.

Love, M.S., C.W. Mecklenburg, T. A. Mecklenburg, and L.K.. Thorsteinson. 2005. *Resource Inventory of Marine and Estuarine Fishes of the West Coast and Alaska: A Checklist of North Pacific and Arctic Ocean Species from Baja California to the Alask-Yukon Border*. U.S. Dept. of Interior, U.S. Geological Survey, Seattle Washington, 276 pp.

Shanks, A.L. and G.L. Eckert. 2005. Population persistence of California Current fishes and benthic crustaceans: a marine drift paradox. *Ecol. Monogr.* 75: 505-524.

Then, A.Y., J.M. Hoenig, N.G. Hall, and D.A. Hewitt. 2015. Evaluating the predictive performance of empirical estimators of natural mortality rate using information on over 200 fish species. *ICES Journal of Marine Science* 72(1): 82–92.

Valle et al. Unpublished data. California Department of Fish and Wildlife Marine Region.

5.2.9 California Halibut

CalCOFI. 2012. Review of selected California fisheries for 2011: Ocean salmon, California sheephead, California halibut, longnose skate, petrale sole, California spiny lobster, Dungeness crab, garibaldi, white shark, and algal blooms, California Department of Fish and Game, Marine Region. Fisheries review CalCOFI Rep., Vol. 53, 2012 http://calcofi.org/publications/calcofireports/v53/Vol_53_Fisheries_15-40.pdf

CDFW. 2011. California halibut stock assessment. Assessors: Maunder, M., P. Reilly, T. Tanaka, G. Schmidt, and K. Penttila. California Department of Fish and Wildlife, Marine Region. <http://www.dfg.ca.gov/marine/sfmp/halibut-assessment.asp>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife. <https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

CDFW. 2015. Commercial Passenger Fishing Vessel Fleet Reported Catches for 2014. California Department of Fish and Wildlife Marine Region. <https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

California Ocean Science Trust (OST). 2013. Rapid Assessments for Selected California Fisheries. August 2013. http://opc.ca.gov/webmaster/ftp/project_pages/Rapid%20Assessments/CA%20Rapid%20Assessments.pdf

Eschmeyer, W.N., E.S. Herald, and H. Hammann. 1983. A field guide to Pacific coast fishes of North America. Houghton Mifflin Company, Boston, U.S.A. 336 pp.

FishBase. *Paralichthys californicus*, California flounder. <http://www.fishbase.org/Summary/speciesSummary.php?ID=4228&AT=California+halibut>

Frey, H.W. (ed.), 1971. California's living marine resources and their utilization. Calif. Dep. Fish Game. 148 pp.

James, K. 2013. Seafood Watch Report California Flounder (California Halibut) *Paralichthys californicus*. U.S. Pacific Bottom Gillnet, Bottom trawl, Hook and line. November 4, 2013. http://www.seafoodwatch.org/-/m/sfw/pdf/reports/f/mba_seafoodwatch_californiaflounder_report.pdf

Jliniak, H.G. 2013. 16 California Halibut, *Paralichthys californicus*. In: Larinto, T. (ed.). Status of the Fisheries Report – An Update Through 2011. California Department of Fish and Wildlife. <https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=65559&inline>

Lesyna, K.M. and C.L Barnes. In Press. Assessment of length- and age-at-maturity for California Halibut (*Paralichthys californicus*), including a histologically-based maturity staging system. California Department of Fish and Wildlife.

MacNair, L.S., M.L. Domeier, and C.S.Y. Chun. 2001. Age, growth, and mortality of California halibut, *Paralichthys californicus*, along southern and central California. Fishery Bulletin. 99: 588-600.
<http://fishbull.noaa.gov/994/mac.pdf>

Maunder, M., P. Reilly, T. Tanaka, G. Schmidt, and K. Penttila. 2011. California Halibut Stock Assessment, Section A: Background information. California Department of Fish and Wildlife.
<https://www.wildlife.ca.gov/Conservation/Marine/NCCFRMP/Halibut-Studies/Halibut-Assessment>

Mondar, M. 2011. California halibut stock assessment: Section B: Assessment of California halibut from the US-Mexico border to Point Conception. Southern California Stock Assessment.
<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=36261>

Reilly, P., T. Tanaka, K. Penttila, G. Schmidt, J. Weinstein, and M. Key. 2008. Applications of Some Recommended Analytical Methods from Data-Poor Workshop (December 2008) to Department of Fish and Game California Halibut Data Sets. <https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=39601>

5.2.10 California Sheephead

Alonzo, S. and M. Mangel. 2004. The effects of size-selective fisheries on the stock dynamics of and sperm limitation in sex-changing fish. Fishery Bulletin 102(1): 1-13.

Alonzo, S. H., M. Key, T. Ish and A. MacCall. 2004. Status of the California sheephead (*Semicossyphus pulcher*) stock. California Department of Fish and Game, Sacramento, California, USA.

Alonzo, S.H., T. Ish, M. Key, A.D. MacCall and M. Mangel M. 2008. The importance of incorporating protogynous sex change into stock assessments. Bull Mar Sci 83: 163–179.

Caselle, J.E., S.L. Hamilton, D.M. Schroeder, M.S. Love, J.D. Standish, J.A. Rosales-Casián and O. Sosa-Nishizaki. 2011. Geographic variation in density, demography, and life history traits of a harvested temperate sex-changing reef fish. Can J. Fish Aquat Sci 68: 288–303.

CDFW. 2013. Status of the Fisheries Report, an Update through 2011. Report to the California Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Wildlife Marine Region. January 2013.
<https://www.wildlife.ca.gov/Conservation/Marine/Status#28027677-status-of-the-fisheries-report-through-2011>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife.
<https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

Cowen, R. K. 1990. Sex change and life history patterns of the labrid, *Semicossyphus pulcher*, across an environmental gradient. Copeia (3): 787-795.

Eschmeyer, W.N., E.S. Herald, and H. Hammann. 1983. A field guide to Pacific coast fishes of North America. Houghton Mifflin Company, Boston, U.S.A. 336 pp.

Hamilton, S.L., J.R. Wilson, T. Ben-Horin, and J.E. Caselle. 2011. Utilizing Spatial Demographic and Life History Variation to Optimize Sustainable Yield of a Temperate Sex-Changing Fish. PLoS ONE 6(9): e24580. doi:10.1371/journal.pone.0024580

Loke, K. 2013. California sheephead. In: Porzio, D. (ed.). Review of selected California fisheries for 2011: Ocean Salmon, California Sheephead, California Halibut, Longnose Skate, Petrale Sole, California Spiny Lobster, Dungeness Crab, Garibaldi, White Shark, and Algal Blooms. CalCOFI Report 53, p. 19-22. http://calcofi.org/publications/calcofireports/v53/Vol_53_Fisheries_15-40.pdf

Love, M. 2011. Certainly More Than You Want to Know About the Fishes of the Pacific Coast. Really Big Press, 672 pp.

MRAG Americas, Inc. 2016. Assessment of CDFW Data Collections, Final Report of Phase 1 & 2. A Report for California Department of Fish and Wildlife and Resources Legacy Fund. MRAG Americas, Essex, MA.

Patrick, W. S., P. Spencer, O. Ormseth, J. Cope, J. Field, D. Kobayashi, T. Gedamke, E. Cortés, K. Bigelow, W. Overholtz, J. Link, and P. Lawson. 2009. Use of productivity and susceptibility indices to determine stock vulnerability, with example applications to six U.S. fisheries. NOAA Tech. Memo. NMFSF/SPO-101, 90 pp.

Sundberg, M.A., K.A. Loke, C.G. Lowe, and K.A. Young. 2009. Gonadal restructuring during sex transition in California sheephead: a reclassification three decades after initial studies. Bull South Calif Acad Sci 108: 16–28.

Topping D.T., C.G. Lowe, and J.E. Caselle. 2006. Site fidelity and seasonal movement patterns of adult California sheephead, *Semicossyphus pulcher* (Labridae), ascertained via longterm acoustic monitoring. Mar Ecol Prog Ser 326: 257-267. <http://www.int-res.com/articles/meps2006/326/m326p257.pdf>

Warner, R. R. 1975. The reproductive biology of the protogynous hermaphrodite *Pimelometopon pulchrum* (Pisces: Labridae). Fishery Bulletin 73: 262-283.

5.2.11 Dungeness Crab

CDFG. 2001. California's Living Marine Resources: A Status Report. The Resources Agency, The California Department of Fish and Game. December 2001. 594 pp. <https://www.wildlife.ca.gov/Conservation/Marine/Status/2001>

CDFW. 2013. Status of the Fisheries Report, an Update through 2011. Report to the California Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Wildlife Marine Region. January 2013. <https://www.wildlife.ca.gov/Conservation/Marine/Status#28027677-status-of-the-fisheries-report-through-2011>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife. <https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

COST. 2013. Rapid Assessments for Selected California Fisheries. August 2013.
http://opc.ca.gov/webmaster/ftp/project_pages/Rapid%20Assessments/CA%20Rapid%20Assessments.pdf

Department of Fisheries and Oceans Canada (DFO). 2013. Pacific Region Integrated Fisheries Management Plan, Crab by Trap, January 1, 2013 to December 31, 2013.

Kashef, N. 2015. Seafood Watch Dungeness crab (*Metacarcinus magister*) British Columbia and United States Pot. October 12, 2015. https://www.seafoodwatch.org/-/m/sfw/pdf/reports/c/mba_seafoodwatch_dungeness_crab_report.pdf

Shanks, A.L. and G.C. Roegner. 2007. Recruitment limitation in Dungeness crab populations is driven by variation in atmospheric forcing. *Ecology*. 88(7): 1726-1737.

Tasto, R.N. 1983. Juvenile Dungeness crab, *Cancer magister*, studies in the San Francisco Bay area. Pages 135-154 in P.W. Wild and R.N. Tasto, eds. Life history, environment, and mariculture studies of the Dungeness crab, *Cancer magister*, with emphasis on the central California fishery resource. Calif. Dep. Fish Game Fish Bull. 172.

Zhang Z., W. Hajas, A. Phillips, and J.A. Boutillier. 2004. Use of length-based models to estimate biological parameters and conduct yield analyses for ale Dungeness crab (*Cancer magister*). *Canadian Journal of Fisheries and Aquatic Sciences* 61: 2126-2134.

5.2.12 Geoduck Clam

CDFG. 2001. California's Living Marine Resources: A Status Report. The Resources Agency, The California Department of Fish and Game. December 2001. 594 pp.
<https://www.wildlife.ca.gov/Conservation/Marine/Status/2001>

MRAG Americas, Inc. 2016. Assessment of CDFW Data Collections - Final Report of Phase 1 & 2. A Report for California Department of Fish and Wildlife and Resources Legacy Fund May 3, 2016.

Trenor, C. and S. Danner. 2008. Seafood Watch Report: Mirugai Pacific Geoduck (*Panopea abrupta*). October 8, 2008. http://www.seafoodwatch.org/-/m/sfw/pdf/reports/g/mba_seafoodwatch_geoduck_report.pdf

5.2.13 Giant Red Sea Cucumber

CDFG. 2008. Status of the Fisheries Report - An Update Through 2006. Report to the Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Game Marine Region, June 2008. <https://www.wildlife.ca.gov/Conservation/Marine/Status#28027679-status-of-the-fisheries-report-through-2006>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife.
<https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

CDFW. 2016. Life History Information for Selected CA Marine Invertebrates and Plants. Accessed August 2016. <https://www.wildlife.ca.gov/Conservation/Marine/Life-History-Inv-And-Plants>

Department of Fisheries and Oceans. 1999. DFO Science Stock Status Report C6-10. Pacific Region Giant Red Sea Cucumber. <http://www.dfo-mpo.gc.ca/csas/Csas/status/1999/C6-10e.pdf>

5.2.14 Jacksmelt

CDFW. 2013. California Marine Sportfish Identification: Other Fishes, Jacksmelt. California Department of Fish and Wildlife Marine Region. <https://www.wildlife.ca.gov/Fishing/Ocean/Fish-ID/Sportfish/Other-Fishes#jacksmelt>

CDFW. 2013. Status of the Fisheries Report, an Update through 2011. Report to the California Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Wildlife Marine Region. January 2013. <https://www.wildlife.ca.gov/Conservation/Marine/Status#28027677-status-of-the-fisheries-report-through-2011>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife. <https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

Clark, F.N. 1929. The Life History of the California Jack Smelt, *Atherinopsis californiensis*. Division of Fish and Game of California, Fish Bulletin No. 16. Contribution No. 77 from the California State Fisheries Laboratory, January, 1929. California State Printing Office, Sacramento. http://content.cdlib.org/view?docId=kt258001x6&brand=calisphere&doc.view=entire_text

Eschmeyer, W.N., E.S. Herald, and H. Hammann. 1983. A field guide to Pacific coast fishes of North America. Houghton Mifflin Company, Boston, U.S.A. 336 pp.

Fishbase. *Atherinopsis californiensis*, Jack silverside. <http://www.fishbase.org/Summary/speciesSummary.php?ID=3236&AT=jacksmelt>

Lavenberg, R.J. and M. Chernoff. 1995. *Atherinidae. Pejerreyes*. p. 889-901. In W. Fischer, F. Krupp, W. Schneider, C. Sommer, K.E. Carpenter and V. Niem (eds.) Guia FAO para Identificación de Especies para lo Fines de la Pesca. Pacifico Centro-Oriental. 3 Vols. FAO, Rome

Shanks, A.L. and G.L. Eckert. 2005. Population persistence of California Current fishes and benthic crustaceans: a marine drift paradox. Ecol. Monogr. 75: 505-524.

Watson, W. 1996. Atherinidae: silversides. p. 608-619. In H.G. Moser (ed.) The early stages of fishes in the California Current region. California Cooperative Oceanic Fisheries Investigations (CalCOFI) Atlas No. 33. Allen Press, Inc., Lawrence, Kansas. 1505 pp.

5.2.15 Kellet's Whelk

CDFG. 2010. California's Living Marine Resources: Status of the Fisheries Report through 2008. Report to the California Fish and Game Commission as directed by the Marine Life Management Act of 1998.

California Department of Fish and Game Marine Region, August 2010.

<https://www.wildlife.ca.gov/Conservation/Marine/Status#28027678-status-of-the-fisheries-report-through-2008>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife.

<https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

Hubbard K. 2008. 2 Kellet's Whelk, *Kelletia kelletii*. [Status of the Fisheries Report](#) 2008 section 2: 1-6.

Romero M. R., K.M. Walker, C.J. Cortez, Y. Sanchez, K.J. Nelson, D.C. Ortega, S.L. Smick, W.J. Hoese, and D.C. Zacherl. 2012. Larval Diel Vertical Migration of the Marine Gastropod *Kelletia kelletii* (Forbes, 1850). Journal of Marine Biology, Article ID 386575, 9 pp. doi:10.1155/2012/386575.

5.2.16 Kelp Bass

CalCOFI Committee. 2009. Review of Selected California Fisheries for 2008: Coastal Pelagic Finfish, Market Squid, Ocean Salmon, Groundfish, California Spiny Lobster, Spot Prawn, White Seabass, Kelp Bass, Thresher Shark, Skates and Rays, Kellet's Whelk and Sea Cucumber. CalCOFI Report, Vol 50, pgs. 30-32. <https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=33769&inline>

CDFW. Unpublished data/ongoing study on survival after capture and release. California Department of Fish and Wildlife Marine Region.

CDFG. 2004. Annual Status of the Fisheries Report Through 2003. Report to the Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Game Marine Region, December 2004.

<https://www.wildlife.ca.gov/Conservation/Marine/Status#28027680-status-of-the-fisheries-report-through-2003>

CDFW. 2013. California Marine Sportfish Identification: Sea Bass. California Department of Fish and Wildlife Marine Region. <https://www.wildlife.ca.gov/Fishing/Ocean/Fish-ID/Sportfish/Sea-Bass#kelp>

CDFW. 2016. Recreational Kelp Bass. California Department of Fish and Wildlife Marine Region. <https://www.wildlife.ca.gov/Conservation/Marine/NCCFRMP/Recreational-Kelp-Bass>

CDFW. 2016. Saltwater Basses Fishery Information. California Department of Fish and Wildlife Marine Region. <https://www.wildlife.ca.gov/Conservation/Marine/SCFRMP/Saltwater-Bass#newregs>

Eschmeyer, W.N., E.S. Herald, and H. Hammann. 1983. A field guide to Pacific coast fishes of North America. Houghton Mifflin Company, Boston, U.S.A. 336 pp.

Fishbase. *Paralabrax clathratus*, Kelp bass.

<http://www.fishbase.org/Summary/speciesSummary.php?ID=3335&AT=kelp+bass>

Heemstra, P.C. 1995. Serranidae. Meros, serranos, guasetas, enjambres, baquetas, indios, loros, gallinas, cabrillas, garropas. p. 1565-1613. In W. Fischer, F. Krupp, W. Schneider, C. Sommer, K.E. Carpenter and

V. Niem (eds.) Guia FAO para Identification de Especies para lo Fines de la Pesca. Pacifico Centro-Oriental. 3 Vols. FAO, Rome.

Jarvis, E.T., H.L. Gliniak, and C.F. Valle. 2014. Effects of fishing and the environment on the long-term sustainability of the recreational saltwater bass fishery in southern California.

<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=90973&inline>

Koengs et al. 2015. Marine Ecology Progress Series. Vol. 537: 1-8. doi: 10.3354/meps11467

Love, M. S., A. Brooks, D. Busatto, J. Stephens, and P. A. Gregory. 1996. Aspects of the life histories of the kelp bass, *Paralabrax clathratus*, and barred sand bass, *P. nebulifer*, from the Southern California Bight. Fishery Bulletin 94: 472-481.

Shanks, A.L. and G.L. Eckert. 2005. Population persistence of California Current fishes and benthic crustaceans: a marine drift paradox. Ecol. Monogr. 75: 505-524.

Then, A.Y., J.M. Hoenig, N.G. Hall, and D.A. Hewitt. 2015. Evaluating the predictive performance of empirical estimators of natural mortality rate information on over 200 fish species. ICES Journal of Marine Science. 72(1): 82-92. doi: 10.1093/icesjms/fsu136.

<http://icesjms.oxfordjournals.org/content/early/2014/08/19/icesjms.fsu136.full.pdf+html>

Young, Parke H. 1963. The Kelp Bass (*Paralabrax clathratus*) and its fishery, 1947-1958. California Fish and Game Bulletin 122: 1-67. <http://content.cdlib.org/ark:/13030/kt609nb14j/>

5.2.17 Market Squid

Baumgartner, T. R., A. Soutar, and V. Ferreira-Bartrina. 1992. Reconstruction of the history of Pacific sardine and Northern anchovy populations over the past two millenia from sediments of the Santa Barbara Basin, California. CalCOFI Reports 33: 24-40.

Butler, J., J. Wagner, and A. Henry. 2001. Age and growth of *Loligo opalescens*. National Marine Fisheries Service Reports.

Butler, J., D. Fuller, and M. Yaremko. 1999. Age and growth of market squid (*Loligo opalescens*) off California during 1998. California Cooperative Oceanic Fisheries Investigations Reports 39: 61-70.

Cailliet, G.M. and D.L. Vaughan. 1983. A Review of the Methods and Problems of Quantitative Assessment of *Loligo opalescens*. Biological Oceanography 2: 379-400.

CDFG. 2008. Status of the Fisheries Report - An Update Through 2006. Report to the Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Game Marine Region, June 2008. <https://www.wildlife.ca.gov/Conservation/Marine/Status#28027679-status-of-the-fisheries-report-through-2006>

CDFW. 2005. Market Squid Fishery Management Plan. California Department of Fish and Wildlife, Marine Region. <https://www.wildlife.ca.gov/Conservation/Marine/MSFMP>

CDFW. 2015. Commercial Passenger Fishing Vessel Fleet Reported Catches for 2014. California Department of Fish and Wildlife, Marine Region.

<https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife.

<https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

COST. 2013. Rapid Assessments for Selected California Fisheries. August 2013.

http://opc.ca.gov/webmaster/ftp/project_pages/Rapid%20Assessments/CA%20Rapid%20Assessments.pdf

Dorval, E., P.R. Crone, and J.D. McDaniel. 2013. Variability of egg escapement, fishing mortality and spawning population in the market squid fishery in the California Current Ecosystem. *Marine and Freshwater Research* 64: 80-90.

Fields, W.G. 1965. The structure, development, food relations, reproduction, and life history of the squid *Loligo opalescens*. California Department of Fish and Game Fish Bulletin 131: 1-108.

Frey, O. 2012. Seafood Watch Report Market Squid *Doryteuthis (Loligo) opalescens*. California Purse Seine. November 9, 2012. http://www.seafoodwatch.org/-/m/sfw/pdf/reports/s/mba_seafoodwatch_camarketsquidreport.pdf

Hurley, A.C. 1977. Mating behavior of the squid *Loligo opalescens*. *Marine Behavior and Physiology* 4(3): 195-203.

Jackson, G. D. 1998. Research into the life history of *Loligo opalescens*: where to from here? *Calif. Coop. Oceanic Fish. Invest. Rep.* 39: 101-107.

Jackson, G.D. and M.L. Domeier. 2003. The effects of an extraordinary El Nino/La Nina event on the size and growth of the squid *Loligo opalescens* off Southern California. *Marine Biology* 142: 925-935.

Macewicz, B.J., J.R. Hunter, N.C.H Lo, and E.L. LaCasella. 2004. Fecundity, egg deposition, and mortality of market squid (*Loligo opalescens*). *Fish Bulletin* 102: 306-327

Morejohn, G.V., J.T. Harvey, and L.T. Krasnow. 1978. The importance of *Loligo opalescens* in the food web of marine vertebrates in Monterey Bay, California. In: C.W. Rechsiek and H.W. Frey. *Biological, oceanographic, and acoustic aspects of the market squid, Loligo opalescens*. Long Beach: California Department of Fish and Game. 67-98.

Protasio, C.Q., A.M. Holder, and B.C. Brady. 2014. Changes in biological characteristics of the California market squid (*Doryteuthis opalescens*) from the California commercial fishery from 2000-01 to 2012-13. *California Fish and Game* 100(2): 276-288.

Reiss, C.S., M.R. Maxwell, and J.R. Hunter. 2004. Investigating environmental effects on population dynamics of *Loligo opalescens* in the Southern California Bight. *CalCOFI Reports* 45: 87-97.

Yang, W.T., R.F. Hixon, P.E. Turk, M.E. Krejci, W.H. Hulet, and R.T. Hanlon. 1986. Growth, behavior, and sexual maturation of the market squid, *Loligo opalescens*, cultured through the life cycle. Fishery Bulletin 84: 771-798.

Zeidberg, L. Year Unknown. *Loligo opalescens*, California Market squid. Accessed August 2016.
<http://www.thecephalopodpage.org/Lopal.php>

5.2.18 Night Smelt

CDFG. 2001. California's Living Marine Resources: A Status Report. The Resources Agency, The California Department of Fish and Game. December 2001. 594 pp.
<https://www.wildlife.ca.gov/Conservation/Marine/Status/2001>

Eschmeyer, W.N., E.S. Herald, and H. Hammann. 1983. A field guide to Pacific coast fishes of North America. Houghton Mifflin Company, Boston, U.S.A. 336 pp.

Fishbase. *Spirinchus starksi*, Night smelt.
<http://www.fishbase.org/Summary/speciesSummary.php?ID=2697&AT=night+smelt>

HT Harvey & Associates. 2015. Collaborative Research on the Spawning Population of Night Smelt (*Spirinchus starksi*) in Humboldt and Del Norte Counties, California. Final Report. HTH Project No. 3501-01. April 15, 2015. <https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=109419&inline>

Moser, H.G. 1996. Osmeridae: smelts. p. 240-243. In H.G. Moser (ed.) The early stages of fishes in the California Current Region. California Cooperative Oceanic Fisheries Investigations (CalCOFI) Atlas No. 33. 1505 pp.

Shanks, A.L. and G.L. Eckert. 2005. Population persistence of California Current fishes and benthic crustaceans: a marine drift paradox. Ecol. Monogr. 75: 505-524.

Therriault et. al. 2002. Review of surf smelt, *Hypomesus pretiosus*, biology and fisheries, with suggested management options for British Columbia.

5.2.19 Ocean Whitefish

Cooksey, D.J. 1980. Age, Growth and Maturity of the Ocean Whitefish, *Caulolatilus princeps*. Thesis.

CDFG. 2004. Annual Status of the Fisheries Report Through 2003. Report to the Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Game Marine Region, December 2004.
<https://www.wildlife.ca.gov/Conservation/Marine/Status#28027680-status-of-the-fisheries-report-through-2003>

Elorduy-Garay, J.F. et al. 2005. Age, growth and mortality of *Caulolatilus princeps* (Pisces: Malacanthidae) from the southern Gulf of California. Hidrobiologica 15 (3): 289-297.

Elorduy-Garay, J.F. and S. Ramirez-Luna. 1994. Gonadal development and spawning of female ocean whitefish, *Caulolatilus princeps* (Pisces: Branchiostegidae) in the Bay of La Paz, B.C.S., Mexico. *Journal of Fish Biology* 44: 553-566.

Eschmeyer, W.N., E.S. Herald, and H. Hammann. 1983. A field guide to Pacific coast fishes of North America. Houghton Mifflin Company, Boston, U.S.A. 336 pp.

Fishbase. *Caulolatilus princeps*, Ocean whitefish.

<http://www.fishbase.org/Summary/SpeciesSummary.php?ID=3539&AT=ocean+whitefish>

Moser GH et al. 1986. Development and Distribution of larvae and pelagic juveniles of Ocean Whitefish, *Caulolatilus princeps*, in the CALCOFI survey region.

5.2.20 Pacific Angel Shark

Cailliet, G.M. 2005. *Squatina californica*. The IUCN Red List of Threatened Species 2005: e.T39328A10202016. <http://dx.doi.org/10.2305/IUCN.UK.2005.RLTS.T39328A10202016.en>.

Cailliet G.M., H.F. Mollet, G.G. Pittenger, D. Bedford, and L.J. Natanson. 1992. Growth and Demography of the Pacific Angel Shark (*Squatina californica*) based upon tag returns off California. *Australian Journal of Marine and Freshwater Research*. 43(5): 1313-1330.

CDFG. 2001. California's Living Marine Resources: A Status Report. The Resources Agency, The California Department of Fish and Game. December 2001. 594 pp.
<https://www.wildlife.ca.gov/Conservation/Marine/Status/2001>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife.
<https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

Compagno, L.J.V. 1984. FAO Species Catalogue. Vol. 4. Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Part 1 - Hexanchiformes to Lamniformes. FAO Fish. Synop. 125(4/1): 1-249. Rome: FAO.

Escobar-Sanchez O., F. Galvan-Magana, and L.A. Abitia-Cardenas. 2011. Trophic level and isotopic composition of d13C and d15N of Pacific Angel Shark, *Squatina californica* (Ayles, 1859), in the Southern Gulf of California, Mexico. *Journal of Fisheries and Aquatic Science*. 6.2: 141-150.

Fishbase. *Squatina californica*, Pacific angelshark.

<http://www.fishbase.org/Summary/speciesSummary.php?ID=729&AT=pacific+angel+shash>

Galvan-Magana F., H.J. Nienhuis, and A.P. Klimley. 1989. Seasonal abundance and feeding habits of sharks of the lower Gulf of California, Mexico. *California Fish and Game*. 75(2): 74-84.

NMFS. 2005. Essential Fish Habitat Designation and Minimization of Adverse Impacts, Pacific Coast Groundfish Fishery Management Plan: Environmental Impact Statement.
http://www.westcoast.fisheries.noaa.gov/publications/nepa/groundfish/final_groundfish_efh_eis.html

Romero-Caicedo AF, F. Galvan-Magana, A. Hernandez-Herrera, and M. Carrera-Fernandez. 2016. Reproductive parameters of the Pacific angel shark *Squatina californica* (Selachii: Squatinidae). Journal of Fish Biology 88(4): 1430-1440.

Smith, S.W., D.W. Au, and C. Show. 1998. Intrinsic rebound potential of 26 species of Pacific sharks. Mar. Freshwat. Res. 49: 663-678.

Standora, E.A. and D.R. Nelson. 1977. A telemetric study of the behavior of free-swimming Pacific angel sharks, *Squatina californica*. Bulletin of the Southern California Academy of Sciences. 76(3): 193-201.

5.2.21 Pacific Hagfish

Barss, W. H. 1993. Pacific hagfish, *Eptatretus stoutii*, and black hagfish, *E. deani*: the Oregon fishery and port sampling observations, 1988-92. Marine Fisheries Review 55(4): 19-30.

CDFG. 2008. Status of the Fisheries Report - An Update Through 2006. Report to the Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Game Marine Region, June 2008. <https://www.wildlife.ca.gov/Conservation/Marine/Status#28027679-status-of-the-fisheries-report-through-2006>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife. <https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

Fishbase. *Eptatretus stoutii*, Pacific hagfish. <http://www.fishbase.org/Summary/speciesSummary.php?ID=2512&AT=pacific+hagfish>

Love, M.S., C.W. Mecklenburg, T.A. Mecklenburg, and L.K. Thorsteinson. 2005. Resource Inventory of Marine and Estuarine Fishes of the West Coast and Alaska: A Checklist of North Pacific and Arctic Ocean Species from Baja California to the Alaska-Yukon Border. U.S. Department of the Interior, U.S. Geological Survey, Biological Resources Division, Seattle, Washington, 98104.

Nakamura, R. 1994. Growth and age of Pacific hagfish *Eptatretus stoutii* off the central California coast. National Oceanic and Atmospheric Administration, National Marine Fisheries Service. Final Report NA27FD0169-01.

5.2.22 Pacific Herring

CDFG. 2008. Status of the Fisheries Report - An Update Through 2006. Report to the Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Game Marine Region, June 2008. <https://www.wildlife.ca.gov/Conservation/Marine/Status#28027679-status-of-the-fisheries-report-through-2006>

CDFW. 2015. Summary of the 2014-15 Pacific Herring Spawning Population and Commercial Fisheries in San Francisco Bay. Final Report. California Department of Fish and Wildlife Marine Region. 17pp. <http://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=108526&inline=true>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife.

<https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

Coad, B.W. and J.D. Reist. 2004. Annotated list of the arctic marine fishes of Canada. Can. MS Rep. Fish Aquat. Sci. 2674:iv:+112 pp.

COST. 2013. Rapid Assessments for Selected California Fisheries. Pacific Herring (*Clupea pallasii*). 14 pp.

http://opc.ca.gov/webmaster/ftp/project_pages/Rapid%20Assessments/Pacific%20Herring.pdf

Fishbase. *Clupea pallasii pallasii*, Pacific herring.

<http://www.fishbase.org/Summary/speciesSummary.php?ID=1520&AT=pacific+herring>

Gunderson, D.R. and P.H. Dygert. 1988. Reproductive effort as a predictor of natural mortality rate. J. Cons. Int. Explor. Mer. 44: 200-209.

Hart, J.L., 1973. Pacific fishes of Canada. Bull. Fish. Res. Board Can. 180. 740 pp.

Lamb, A. and P. Edgell. 1986. Coastal fishes of the Pacific northwest. Harbour Publishing Co. Ltd., B.C., Canada. 224 pp. DOI <http://dx.doi.org/>

Matarese, A.C., A.W. Kendall, D.M. Blood, and M.V. Vinter. 1989. Laboratory guide to early life history stages of Northeast Pacific fishes. NOAA Tech. Rep. NMFS 80: 1-652.

Morrow, J.E. 1980. The freshwater fishes of Alaska. University of B.C. Animal Resources Ecology Library. 248 pp.

MRAG Americas, Inc. 2016. Assessment of CDFW Data Collections - Final Report of Phase 1 & 2. A Report for California Department of Fish and Wildlife and Resources Legacy Fund May 3, 2016.

Musick, J.A., M.M. Harbin, S.A. Berkeley, G.H. Burgess, A.M. Eklund, L. Findley, R.G. Gilmore, J.T. Golden, D.S. Ha, G.R. Huntsman, J.C. McGovern, S.J. Parker, S.G. Poss, E. Sala, T.W. Schmidt, G.R. Sedberry, H. Weeks, and S.G. Wright, 2000. Marine, estuarine, and diadromous fish stocks at risk of extinction in North America (exclusive of Pacific salmonids). Fisheries 25(11): 6-30.

Schaefer, M.B., 1937. Notes on the spawning of the Pacific herring *Clupea pallasii*. Copeia 1937(1): 57.

Siple, M. 2014. Seafood Watch Report Pacific herring *Clupea pallasii*. California Bottom Gillnet. August 21, 2014. <http://www.seafoodwatch.org> -

http://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=108526&inline=true/m/sfw/pdf/reports/h/mba_s_eafoodwatch_pacificherring_california_report.pdf

Trumble, R.J. and R.D. Humphreys. 1985. Management of Pacific herring (*Clupea harengus pallasii*) in the eastern Pacific Ocean. Can. J. Fish. Aquat. Sci. 42 (Suppl. 1): 230-244.

5.2.23 Pink Shrimp

CDFG. 2008. Status of the Fisheries Report - An Update Through 2006. Report to the Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Game Marine Region, June 2008. <https://www.wildlife.ca.gov/Conservation/Marine/Status#28027679-status-of-the-fisheries-report-through-2006>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife. <https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

COST. 2013. Rapid Assessments for Selected California Fisheries. http://opc.ca.gov/webmaster/ftp/project_pages/Rapid%20Assessments/CA%20Rapid%20Assessments.pdf

Dahlstrom, WA. FAO Fisheries Synopsis No. 99. Synopsis of Biological Data on the Ocean Shrimp. <http://www.fao.org/docrep/005/ac765t/ac765t11.htm>

Intertek Moody Marine. 2013. Oregon Pink Shrimp (*Pandalus jordani*) Trawl Fishery. PUBLIC CERTIFICATION REPORT. February 12, 2013. https://www.msc.org/track-a-fishery/fisheries-in-the-program/certified/pacific/oregon-and-washington-pink-shrimp/reassessment-downloads-1/20130214_PCR_SHR94.pdf

MRAG Americas, Inc. 2016. Assessment of CDFW Data Collections - Final Report of Phase 1 & 2. A Report for California Department of Fish and Wildlife and Resources Legacy Fund May 3, 2016.

Townsend, S. 2014. Seafood Watch: Coonstripe shrimp, Dock shrimp, pink shrimp, Northern shrimp, Sidesripe shrimp, Spot prawn, Ridgeback shrimp. Feb 2014. http://www.seafoodwatch.org/-/m/sfw/pdf/reports/s/mba_seafoodwatch_coldwatershrimp_westcoastus_report.pdf

5.2.24 Pismo Clam

CDFG. 2008. Status of the Fisheries Report - An Update Through 2006. Report to the Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Game Marine Region, June 2008. <https://www.wildlife.ca.gov/Conservation/Marine/Status#28027679-status-of-the-fisheries-report-through-2006>

5.2.25 Red Abalone

Allee, W. C. 1931. Co-operation among animals. Am. J. Sociol. 37: 386–398.

Babcock, R. and J. Keesing. 1999. Fertilization biology of the abalone *Haliotis laevis*: laboratory and field studies. Can. J. Fish. Aquat. Sci. 56: 1668–1678.

Button, C. A. 2008. The influence of density-dependent aggregation characteristics on the population biology of benthic broadcast spawning gastropods: pink abalone (*Haliotis corrugata*), red abalone (*Haliotis rufescens*), and wavy turban snails *Megastrea undosa*. PhD diss., University of California at San Diego. 209 pp.

CDFG. 2001. Draft supplement to the environmental document - ocean sport fishing regulations concerning abalone sport fishing. The Resources Agency, Sacramento, California.

CDFW. 2010. Abalone Recovery and Management Plan Status Report - Northern California Red Abalone Fishery Marine Region Invertebrate Management Project. May 2010.

<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=29511&inline>

CDFW. 2013. Estimating Red Abalone Density for Managing California's Recreational Red Abalone Fishery. October 2013. <https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=123709&inline>

CDFW. 2013. Fishery Status Update: Northern California Red Abalone (2009-2012).

<https://www.wildlife.ca.gov/Conservation/Marine/Invertebrates/Abalone/Abalone-Status>

CDFW. 2013. Status of the Fisheries Report, an Update through 2011. Report to the California Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Wildlife Marine Region. January 2013.

<https://www.wildlife.ca.gov/Conservation/Marine/Status#28027677-status-of-the-fisheries-report-through-2011>

CDFW. 2016. Invertebrates of Interest: Abalone. California Department of Fish and Wildlife Marine Region. <https://www.wildlife.ca.gov/Conservation/Marine/Invertebrates/Abalone#29972974-how-do-abalone-reproduce>

CDFW. Red Abalone FAQs.

<https://www.wildlife.ca.gov/Conservation/Marine/Invertebrates/Abalone#29972979-are-abalone-vulnerable-to-overfishing>

Leaf, R. T., L. Rogers-Bennett, and P. L. Haaker. 2007. Spatial, temporal, and size-specific variation in mortality estimates of red abalone, *Haliotis rufescens*, from mark-recapture data in California. Fisheries Research 83: 341–350.

Rogers-Bennett, L., R.F. Dondanville, and J. Kashiwada. 2004. Size Specific Fecundity of Red Abalone (*Haliotis rufescens*): Evidence for Reproductive Senescence. Journal of Shellfish Research. Vol. 23. No. 2. 553-560. <https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=121986>

Rogers-Bennet, L., D.W. Rogers, and S.A. Schultz. 2007. Modeling Growth and Mortality of Red Abalone (*Haliotis rufescens*). Journal of Shellfish Research 26(3): 719-727.

5.2.26 Red Sea Urchin

CDFG. 2001. California's Living Marine Resources: A Status Report. The Resources Agency, The California Department of Fish and Game. December 2001. 594 pp.

<https://www.wildlife.ca.gov/Conservation/Marine/Status/2001>

CDFG. 2004. Annual Status of the Fisheries Report Through 2003. Report to the Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Game Marine Region, December 2004.

<https://www.wildlife.ca.gov/Conservation/Marine/Status#28027680-status-of-the-fisheries-report-through-2003>

CDFW. 2015. Final California Commercial Landings for 2014. Table 15. California Department of Fish and Wildlife Marine Region. <https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

Lopuch, M. 2008. Seafood Watch Report. Red Sea Urchin (*Strongylocentrotus franciscanus*), Green Sea Urchin (*Strongylocentrotus droebachiensis*). Final Report. September 3, 2008.
http://www.seafoodwatch.org/-/m/sfw/pdf/reports/u/mba_seafoodwatch_urchinreport.pdf

Fisheries and Oceans Canada (DFO). 2000. Underwater World Red Sea Urchin. Communication Directorate Fisheries and Oceans Canada, Ottawa, Ontario.

5.2.27 Redtail Surfperch

Breder, C.M. and D.E. Rosen. 1966. Modes of reproduction in fishes. T.F.H. Publications, Neptune City, New Jersey. 941 pp.

CDFW. 2013. Status of the Fisheries Report, an Update through 2011. Report to the California Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Wildlife Marine Region. January 2013.
<https://www.wildlife.ca.gov/Conservation/Marine/Status#28027677-status-of-the-fisheries-report-through-2011>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife.
<https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

CDFW. 2016. Current California Ocean Recreational Fishing Regulations. California Department of Fish and Wildlife Marine Region. <https://www.wildlife.ca.gov/Fishing/Ocean/Regulations/Fishing-Map/Mendocino>

CDFW. Unpublished data/ongoing studies on growth and natural mortality. California Department of Fish and Wildlife Marine Region.

Eschmeyer, W.N., E.S. Herald, and H. Hammann. 1983. A field guide to Pacific coast fishes of North America. Houghton Mifflin Company, Boston, U.S.A. 336 pp.

Fishbase. *Amphistichus rhodotus*, Redtail surfperch.
<http://www.fishbase.org/Summary/speciesSummary.php?ID=3624&AT=redtail+surfperch>

Shanks, A.L. and G.L. Eckert. 2005. Population persistence of California Current fishes and benthic crustaceans: a marine drift paradox. Ecol. Monogr. 75: 505-524.

5.2.28 Ridgeback Prawn

Allen, M.J., D. Cadien, E. Miller, D.W. Diehl, K. Ritter, S.L. Moore, C. Cash, D.J. Pondella, V. Raco-Rands, C. Thomas, R. Gartman, W. Power, A.K. Latker, J. Williams, J. L. Armstrong, and K. Schiff. 2011. Southern

California Bight 2008 Regional Monitoring Program: Volume IV. Demersal Fishes and Megabenthic Invertebrates. Southern California Coastal Water Research Project, Costa Mesa, CA.

http://ftp.sccwrp.org/pub/download/DOCUMENTS/TechnicalReports/655_B08Trawl.pdf

Anderson, S.L., L.W. Botsford, and W.H. Clark Jr. 1985. Size Distributions and Sex Ratios of Ridgeback prawn (*Sicyonia ingentis*) in the Santa Barbara Channel (1979-1981). CalCOFI. Vol. 26.

http://www.calcofi.org/publications/calcofireports/v26/Vol_26_Anderson_etal.pdf

CDFG. 2008. Status of the Fisheries Report - An Update Through 2006. Report to the Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Game Marine Region, June 2008. <https://www.wildlife.ca.gov/Conservation/Marine/Status#28027679-status-of-the-fisheries-report-through-2006>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife.

<https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

MRAG Americas, Inc. 2016. Assessment of CDFW Data Collections - Final Report of Phase 1 & 2. A Report for California Department of Fish and Wildlife and Resources Legacy Fund. May 3, 2016.

NMFS. 2005. Essential Fish Habitat Designation and Minimization of Adverse Impacts, Pacific Coast Groundfish Fishery Management Plan: Environmental Impact Statement.

http://www.westcoast.fisheries.noaa.gov/publications/nepa/groundfish/final_groundfish_efh_eis.html

Townsend, S. 2014. Seafood Watch: Coonstripe shrimp, Dock shrimp, pink shrimp, Northern shrimp, Sidesripe shrimp, Spot prawn, Ridgeback shrimp. Feb 2014. http://www.seafoodwatch.org/-/m/sfw/pdf/reports/s/mba_seafoodwatch_coldwatershrimp_westcoastus_report.pdf

5.2.29 Shiner Perch

Breder, C.M. and D.E. Rosen. 1966. Modes of reproduction in fishes. T.F.H. Publications, Neptune City, New Jersey. 941 pp.

CDFW. 2013. Status of the Fisheries Report, an Update through 2011. Report to the California Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Wildlife Marine Region. January 2013.

<https://www.wildlife.ca.gov/Conservation/Marine/Status#28027677-status-of-the-fisheries-report-through-2011>

Eschmeyer, W.N., E.S. Herald, and H. Hammann. 1983. A field guide to Pacific coast fishes of North America. Houghton Mifflin Company, Boston, U.S.A. 336 pp.

Fishbase. *Cymatogaster aggregate*, Shiner perch.

<http://www.fishbase.org/Summary/speciesSummary.php?ID=3626&AT=shiner+perch>

Morrow, J.E. 1980. The freshwater fishes of Alaska. University of B.C. Animal Resources Ecology Library. 248 pp.

Shanks, A.L. and G.L. Eckert. 2005. Population persistence of California Current fishes and benthic crustaceans: a marine drift paradox. *Ecol. Monogr.* 75: 505-524.

5.2.30 Spiny Lobster

CDFG. 2001. California's Living Marine Resources: A Status Report. The Resources Agency, The California Department of Fish and Game. December 2001. 594 pp.

<https://www.wildlife.ca.gov/Conservation/Marine/Status/2001>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife.

<https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

CDFW. 2016. California Spiny Lobster Fishery Management Plan. April, 2016. California Department of Fish and Wildlife Marine Region. 239 pp.

<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=121938&inline>

CDFW and CA Sea Grant. 2015. California Spiny Lobster Fishing and Life History Information. California Department of Fish and Wildlife Marine Region.

<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=36321&inline>

Neilson, D.J. 2011. Assessment of the California Spiny Lobster (*Panulirus interruptus*). California Department of Fish and Game, San Diego, CA.

<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=41068&inline>

5.2.31 Spot Prawn

CDFG. 2008. Status of the Fisheries Report - An Update Through 2006. Report to the Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Game Marine Region, June 2008. <https://www.wildlife.ca.gov/Conservation/Marine/Status#28027679-status-of-the-fisheries-report-through-2006>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife.

<https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

MRAG Americas, Inc. 2016. Assessment of CDFW Data Collections - Final Report of Phase 1 & 2. A Report for California Department of Fish and Wildlife and Resources Legacy Fund. May 3, 2016.

NMFS. 2005. Essential Fish Habitat Designation and Minimization of Adverse Impacts, Pacific Coast Groundfish Fishery Management Plan: Environmental Impact Statement.

http://www.westcoast.fisheries.noaa.gov/publications/nepa/groundfish/final_groundfish_efh_eis.html

NOAA. 2011. Voices of the Bay: Fishery Basics – California Fisheries Spot Prawn (*Pandalus platyceros*) . December 2011. <http://sanctuaries.noaa.gov/education/voicesofthebay/pdfs/spotprawns.pdf>

Townsend, S. 2014. Seafood Watch: Coonstripe shrimp, Dock shrimp, pink shrimp, Northern shrimp, Sidedstripe shrimp, Spot prawn, Ridgeback shrimp. Feb 2014. http://www.seafoodwatch.org/-/m/sfw/pdf/reports/s/mba_seafoodwatch_coldwatershrimp_westcoastus_report.pdf

5.2.32 Spotted Sand Bass

Allen et al. 1995. The life history of the spotted sand bass (*Paralabrax maculatofasciatus*) within the Southern California Bight. California Cooperative Oceanic Fisheries Investigations Reports. 36: 196-203.

Anonymous. 1994. Atlas pesquero de México. Instituto Nacional de la Pesca. 234 pp.

CDFG. 2004. Annual Status of the Fisheries Report Through 2003. Report to the Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and Game Marine Region, December 2004.

<https://www.wildlife.ca.gov/Conservation/Marine/Status#28027680-status-of-the-fisheries-report-through-2003>

CDFW. 2013. California Marine Sportfish Identification: Sea Bass. California Department of Fish and Wildlife Marine Region. <https://www.wildlife.ca.gov/Fishing/Ocean/Fish-ID/Sportfish/Sea-Bass#kelp>

CDFW. 2016. Saltwater Basses Fishery Information. California Department of Fish and Wildlife Marine Region. <https://www.wildlife.ca.gov/Conservation/Marine/SCFRMP/Saltwater-Bass#newregs>

de Mitcheson, Y.S. and M. Liu. 2008. Functional hermaphroditism in teleosts. Fish Fish. 9: 1-43.

Fishbase. *Paralabrax maculatofasciatus*, Spotted sand bass.

<http://www.fishbase.org/Summary/speciesSummary.php?ID=3336&AT=spotted+sand+bass>

Heemstra, P.C. 1995. Serranidae. Meros, serranos, guasetas, enjambres, baquetas, indios, loros, gallinas, cabrillas, garropas. p. 1565-1613. In W. Fischer, F. Krupp, W. Schneider, C. Sommer, K.E. Carpenter, and V. Niem (eds.) Guia FAO para Identificación de Especies para lo Fines de la Pesca. Pacifico Centro-Oriental. 3 Vols. FAO, Rome.

Jarvis, E.T., H.L. Gliniak, and C.F. Valle. 2014. Effects of fishing and the environment on the long-term sustainability of the recreational saltwater bass fishery in southern California.

<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=90973&inline>

Shanks, A.L. and G.L. Eckert. 2005. Population persistence of California Current fishes and benthic crustaceans: a marine drift paradox. Ecol. Monogr. 75: 505-524.

Then, A.Y., J.M. Hoenig, N.G. Hall, and D.A. Hewitt. 2015. Evaluating the predictive performance of empirical estimators of natural mortality rate using information on over 200 fish species. ICES Journal of Marine Science 72(1): 82-92.

5.2.33 Warty Sea Cucumber

CDFG. 2008. Status of the Fisheries Report - An Update Through 2006. Report to the Fish and Game Commission as directed by the Marine Life Management Act of 1998. California Department of Fish and

Game Marine Region, June 2008. <https://www.wildlife.ca.gov/Conservation/Marine/Status#28027679-status-of-the-fisheries-report-through-2006>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife.
<https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

CDFW. 2016. Life History Information for Selected CA Marine Invertebrates and Plants. Accessed August 2016. <https://www.wildlife.ca.gov/Conservation/Marine/Life-History-Inv-And-Plants>

Department of Fisheries and Oceans Canada. 1999. DFO Science Stock Status Report C6-10. Pacific Region Giant Red Sea Cucumber. <http://www.dfo-mpo.gc.ca/csas/Csas/status/1999/C6-10e.pdf>

Sanctuary Integrated Monitoring Network. 2016. Species Database, *Parastichopus parvimensis* - Warty sea cucumber. Accessed August 2016.
<http://sanctuarymonitoring.org/species/parastichopus/parvimensis/warty-sea-cucumber>

5.2.34 White Croaker

CDFW. 2013. California Marine Sportfish Identification: Croakers. California Department of Fish and Wildlife Marine Region. <https://www.wildlife.ca.gov/Fishing/Ocean/Fish-ID/Sportfish/Croakers#white>

Eschmeyer, W.N., E.S. Herald, and H. Hammann. 1983. A field guide to Pacific coast fishes of North America. Houghton Mifflin Company, Boston, U.S.A. 336 pp.

FishBase. *Genyonemus lineatus*, White croaker.
<http://www.fishbase.org/Summary/SpeciesSummary.php?ID=422&AT=white+croaker>

Frey, H.W. (ed.). 1971. California's living marine resources and their utilization. Calif. Dep. Fish Game. 148 pp.

Louie, A. and H.G. Gliniak. 2013. White croaker, *Genyonemus lineatus*. In: Larinto, T. (ed.). Status of the Fisheries Report – An Update Through 2011. CA Dept. of Fish and Wildlife.
<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=65553&inline>

Love, M.S., G.E. McGowen, W. Westphal, R.J. Lavenberg, and L. Martin. 1984. Aspects of the life history and fishery of the white croaker, *Genyonemus lineatus* (Sciaenidae), off California. Fish. Bull. 82: 179-198.

Macchi, G.J. 1998. Preliminary estimate of spawning frequency and batch fecundity of striped weakfish, *Cynoscion striatus*, in coastal waters off Buenos Aires province. Fish. Bull. 96(2): 375-381.

Moore, S.L. 1998. Age and growth of white croaker (*Genyonemus lineatus*) off Palos Veredes and Dana Point, California.

Shanks, A.L. and G.L. Eckert. 2005. Population persistence of California Current fishes and benthic crustaceans: a marine drift paradox. Ecol. Monogr. 75: 505-524.

Wolfe, B.W. and C.G. Lowe. 2015. Movement patterns, habitat use and site fidelity of the white croaker in the Palos Verdes Superfund Site, Los Angeles, California. Marine Environmental Research.

5.2.35 White Seabass

Aalbers, S. A. and C. A. Sepulveda. 2015. Seasonal movement patterns and temperature profiles of adult white seabass *Atractoscion nobilis* off of California. Fish Bull. 113: 1–14.

Allen, L., D. Pondella, and M. Shane. 2007. Fisheries Independent Assessment of a Returning Fishery: Abundance of Juvenile White Seabass (*Atractoscion nobilis*) in the Shallow Nearshore Waters of the Southern California Bight, 1995-2005. Fisheries Research. 88: 24-32.

CA Fisheries Fund. 2016. California Fisheries Atlas - White Seabass. (Accessed June 2016). California Department of Fish and Wildlife Marine Region.

http://www.californiafisheriesfund.org/reso_atlas_sbass.html

CDFG. 2002. Final White Seabass Fishery Management Plan. California Department of Fish and Game.

<https://www.wildlife.ca.gov/Conservation/Marine/WSFMP>

CDFW. 2010. White Seabass Enhancement Plan. June 2010. California Department of Fish and Wildlife Marine Region.

<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=29458>

CDFW. 2015. Final California Commercial Landings for 2014 (Table 15). State of California, The Natural Resources Agency, Department of Fish and Wildlife.

<https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

CDFW. 2015. Commercial Passenger Fishing Vessel Fleet Reported Catches for 2014. California Department of Fish and Wildlife Marine Region.

<https://www.wildlife.ca.gov/Fishing/Commercial/Landings#26004609-2014>

CDFW. 2015. White Seabass Fishery Management Plan 2013-2014 Annual Review April 2015. California Department of Fish and Wildlife Marine Region.

<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=100521&inline>

CDFW. 2016. Ocean Resources Enhancement and Hatchery Program Webpage. Accessed August 2016. California Department of Fish and Wildlife Marine Region.

<https://www.wildlife.ca.gov/Conservation/Marine/ABMP/OREHP>

CDFW. 2016. Recreational regulations (Accessed June 2016). California Department of Fish and Wildlife Marine Region. <https://www.wildlife.ca.gov/Fishing/Ocean/Regulations/Fishing-Map/southern>

COST. 2013. Rapid Assessments for Selected California Fisheries. August 2013.

http://opc.ca.gov/webmaster/ftp/project_pages/Rapid%20Assessments/CA%20Rapid%20Assessments.pdf

- Donohoe, C.J. 1997. Age, growth, distribution, and food habits of recently settled white seabass, *Atractoscion nobilis*, off San Diego country, California. Fish. Bull. 95: 709-721.
- Eschmeyer, W.N., E.S. Herald, and H. Hammann. 1983. A field guide to Pacific coast fishes of North America. Houghton Mifflin Company, Boston, U.S.A. 336 pp.
- Franklin, M.P. 1997. An investigation into the population structure of white seabass (*Atractoscion nobilis*), in California and Mexican waters using microsatellite DNA analysis. PhD. Dissertation. University of California Santa Barbara. 109 pp.
- Franklin, M. P., C.L. Chabot, and L.G. Allen. 2016. A Baseline Investigation into the Population Structure of White Seabass, *Atractoscion nobilis*, in California and Mexican Waters Using Microsatellite DNA Analysis. Bull. Southern California Acad. Sci. 115(2): 126–135
- James, K. 2014. MBA Seafood Watch Report California yellowtail, White seabass (*Seriola lalandi*, *Atractoscion nobilis*). California Bottom gillnet, Drift gillnet, Hook and Line. February 13, 2014. http://www.seafoodwatch.org/-/m/sfw/pdf/reports/s/mba_seafoodwatch_cayellowtail_whitesebassreport.pdf
- Karpov, K.A., D.P. Albin, and W.H. Van Buskirk. 1995. The marine recreational fishery in northern and central California, a historical comparison (1958-86), status of stocks (1980-86), and effects of changes in the California current. Calif. Dept. Fish Game, Fish Bull. 176: 1-192.
- Love, M. 1996. Probably more than you want to know about the fishes of the Pacific coast. Really Big Press. Santa Barbara, California.
- Radovich, J. 1961. Relationships of some marine organisms of the northeast Pacific to water temperatures, particularly during 1957 through 1959. Calif. Dept. Fish Game, Fish Bull. 54: 1- 62.
- Romo-Curiel, A. E., S.Z. Herzka, O. Sosa-Nishizaki, C.A. Sepulveda, and S. A. Aalbers. 2015. Otolith-based growth estimates and insights into population structure of White Seabass, *Atractoscion nobilis*, off the Pacific coast of North America. Fisheries Research. 161: 374-383.
- Shanks, A.L. and G.L. Eckert. 2005. Population persistence of California Current fishes and benthic crustaceans: a marine drift paradox. Ecol. Monogr. 75: 505-524.
- Skogsberg, T. 1939. The fishes of the family Sciaenidae (croakers) of California. Calif. Div. Fish Game, Fish Bull. 54: 1- 62.
- Stevens, M., J. Bausell, B. Botkin, and W. Norden. 2011. Seafood Watch Seafood Report, White Seabass (*Atractoscion nobilis*) California. January 10, 2011. http://www.seachoice.org/wp-content/uploads/2011/12/MBA_SeafoodWatch_WhiteSeabassReport.pdf
- Thomas, J. 1968. Management of the white seabass (*Cynoscion nobilis*) in California waters. Calif. Fish. Game Bull. 142. 34 pp.

Valero, J. and L. Waterhouse. 2016. California White Seabass Stock Assessment in 2016.
http://www.capamresearch.org/sites/default/files/WSB_SA_Report_2016.pdf

5.2.36 White Sturgeon

CDFG. 2001. California's Living Marine Resources: A Status Report. The Resources Agency, The California Department of Fish and Game. December 2001. 594 pp.
<https://www.wildlife.ca.gov/Conservation/Marine/Status/2001>

CDFW. 2010. WHITE STURGEON *Acipenser transmontanus* (Richardson). California Department of Fish and Wildlife Marine Region. <https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=104390>

CDFW. 2016. Recreational Fishing Regulations. Accessed Aug 2016. California Department of Fish and Wildlife Marine Region. <https://www.wildlife.ca.gov/Fishing/Ocean/Regulations/Fishing-Map/Central>

CDFW. Unpublished data. California Department of Fish and Wildlife Marine Program.

DuBois, J., M. Gingras, and G. Aasen. 2011. Status and Trends of San Francisco Estuary White Sturgeon. Interagency Ecological Program for the San Francisco Estuary (IEP) Newsletter 24(1): 50.
<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentId=44050>

DuBois, J. and M.D. Harris. 2016. 2015 Sturgeon Fishing Report Card: Preliminary Data Report. California Department of Fish and Wildlife Bay Delta Region.
<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentId=121293>

Fish, M.A. 2010. A White Sturgeon Year-Class Index for the San Francisco Estuary and Its Relation to Delta Outflow. Interagency Ecological Program for the San Francisco Estuary (IEP) Newsletter 23(2): 80.
<https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentId=26542>

Fishbase. *Acipenser transmontanus*, White sturgeon.
<http://www.fishbase.org/Summary/speciesSummary.php?ID=2594&AT=white+sturgeon>

Morrow, J.E. 1980. The freshwater fishes of Alaska. University of. B.C. Animal Resources Ecology Library. 248 pp.

Rien, T.A. and R.C. Beamesderfer. 1994. Accuracy and precision of white sturgeon age estimates from pectoral fin rays. Trans. Am. Fish. Soc. 123(2): 255-265.

Scott, W.B. and E.J. Crossman. 1973. Freshwater fishes of Canada. Bull. Fish. Res. Board Can. 184: 1-966.